

Identification of Bioactive Diterpenoid Lactones using Nuclear Magnetic Resonance from *Andrographis paniculata*

Raghava N. Sriramaneni^{1,2}, Amirin Sadikun², Naveen Kumar Hawala Shivashekaregowda³, Annegowda Venkatappa⁴, Mohamad Zaini Asmawi²

¹Department of Human Oncology, Wisconsin Institute for Medical Research, University of Wisconsin, Wisconsin, Madison, USA, ²Department of Pharmacology, School of Pharmaceutical Sciences, Universiti Sains Malaysia, Pulau Pinang, ³Department of Pharmacology, School of Pharmacy, Taylor's University, Kula Lumpur, Malaysia, ⁴Department of Phytochemistry, Adhichunchangiri University, Mandya, Karnataka, India

ABSTRACT


Background: *Andrographis paniculata* (AP) (Burm. F) Nees, belongs to the family *Acanthaceae*, it is also known as a highly bitter plant. In South East Asia this plant is commonly used for cold, cough and respiratory tract infections. Andrographolide (ANG), 14-deoxy-11, 12-didehydroandrographolide (DDA) are the major bioactive diterpenoid lactone compounds of the plant shown vasorelaxant effect, anticancer, anti-inflammatory, and to treat variety of diseases which is well-documented. **Materials and Methods:** The high-performance liquid chromatography (HPLC) and ¹H-NMR analysis of AP chloroform extract (APCE) revealed the presence of ANG, 14-deoxyandrographolide (DA), and DDA. The main advantage of ¹H-NMR is simple, rapid and successfully applied to quantify the active diterpenoids and an alternative to HPLC method to check the presence of various bioactive compounds and its presence in the active raw materials of AP. **Results:** The results revealed that most active bioactive compounds such as ANG, DA, and DDA are present in the APCE portion of the AP. The chemical shifts of various groups confirmed through ¹H-NMR that all the three ANG, DA, and DDA has potent vasorelaxant effect and majority of these compounds are present in the APCE portion and not in the methanol or aqueous fractions. The ¹H-NMR study confirmed that diterpenoids such as ANG, DA, and DDA from APCE has vasorelaxant effect on the animal study.

Key words: ¹H-NMR, *Andrographis paniculata*, andrographolide, bioactive compounds, diterpenoids, vasorelaxant

SUMMARY

In this present study, an ¹H-NMR method was successfully developed to quantify four major diterpenoids (andrographolide, dehydroandrographolide, deoxyandrographolide, and neoandrographolide) in *Andrographis paniculata* chloroform extract and its commercial preparations by using a novel solvent system containing CDCl₃. This analytical method was proved to be of excellent accuracy, precision, and repeatability, thus it is simply affordable, rapid,

and powerful tool for efficient quality assessment. For further validation, the total content and determination of two diterpenoids 14-deoxy andrographolide and 14-deoxy-11, 12-didehydroandrographolide in *Andrographis paniculata* chloroform extract were accomplished. The results were in good contrast with those determined by high-performance liquid chromatography method. Compared with high-performance liquid chromatography, ¹H-NMR method is simple, rapid, and reliable without the creation of calibration curves, significantly much short analysis time, and thus is more suitable for the routine control of *Andrographis paniculata* chloroform extract and its viable preparations.



Abbreviations Used: ANG: Andrographolide; APCE: *Andrographis paniculata* chloroform extract; DA: 14-deoxy andrographolide; DDA: 14-deoxy-11, 12-didehydroandrographolide; AP: *Andrographis paniculata*.

Correspondence:

Dr. Raghava N. Sriramaneni,
Department of Human Oncology, Wisconsin
Institute for Medical Research, Wisconsin, USA.
E-mail: sriramaneni@wisc.edu
DOI: 10.4103/pr_117_18

Access this article online

Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

Andrographolide (ANG), dehydroandrographolide, deoxyandrographolide, and neoandrographolide are the four most dominant diterpenoids, proven for a great proportion of components in *Andrographis paniculata* (AP), showing diverse activities, for example, anti-inflammatory, antiviral, immune-stimulatory, and hepatoprotective activities both *in vitro* and *in vivo* experimental studies.^[1-4]

Structures of four major bioactive diterpenoids from AP, highlighted lines are the structural differences are noticed. However, various influences such as geographic region, crop time, and processing method lead to the variability in the content of these active diterpenoids.^[5,6] Hence, it is very crucial to develop a practical analysis method for both the raw materials and isolated compounds of the natural plants. Various chromatographic methods such as thin layer chromatography, high-performance liquid chromatography, and column chromatography and capillary electrophoresis have already reported. In addition to that, these methods

are time-consuming, not sensitive and most complicated methods when compared to ¹H-NMR analysis.

Nuclear magnetic resonance (NMR) spectroscopy is the study of molecules by recording the interaction of radiofrequency electromagnetic radiation with the nuclei of molecules placed in a strong magnetic field.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Sriramaneni RN, Sadikun A, Shivashekaregowda NK, Venkatappa A, Asmawi MZ. Identification of bioactive diterpenoid lactones using nuclear magnetic resonance from *Andrographis paniculata*. Phcog Res 2019;11:121-6.




Figure 1: Chemical structure of Andrographolide. Molecular formula: $C_{20}H_{30}O_5$. Molecular weight: 350.46 g

Like all other spectroscopic techniques, NMR spectroscopy involves the interaction of the material being examined with electromagnetic radiation.

NMR spectroscopy is a useful technique for identifying and analyzing organic compounds. This extremely important experimental technique is based on the magnetic nuclear spin of 1H , ^{13}C , ^{15}N , ^{19}F , ^{31}P , and so forth. Only 1H and ^{13}C will give more detailed and definitive information.

Hence, when information obtained from NMR and other analytical methods are compared, it is apparent that the most detailed and necessary information to recognize a structure is obtained from NMR spectra. However, this does not mean that NMR is always sufficient for structural analysis. For absolute analysis of structures, it is necessary to apply all spectroscopic methods. If a new compound is synthesized, the chemist who does scientific research should have all the spectroscopic data related to this chemical compound. The information provided by an NMR spectrum is useful only if it can be interpreted well. In addition to providing information related to constitutional analysis, NMR spectra provide detailed information pertinent to configurational and conformational analysis.^[7]

The aerial parts of the plant AP (leaves and stems) are used to extract the active phytochemicals. Previous investigations on the chemical composition of AP showed that it is a rich source of diterpenoids and 2'-oxygenated flavonoids including ANG, neoandrographolide, 14-deoxy-11, 12-didehydroandrographolide (DDA), 14-deoxyandrographolide (DA), isoandrographolide, and DA-19 β -D-glucoside, homoandrographolide, andrographan, andrographasterin, and stigmasterol.^[8]

The primary bioactive diterpenoid lactone compound of the medicinal plant AP is andrographolide (ANG).^[9-11] The chemical formula is $[C_{20}H_{30}O_5]$: 3-(2-{decahydro-6-hydroxy-5-(hydroxymethyl)-5, 8 α -dimethyl-2-methylene-1-pthalenyl} thyledene) dihydro-4-hydroxy-2 (3H)-furanone; Figure 1 is a colorless crystalline bicyclic diterpenoid lactone and has a very bitter taste.^[12] Based on previous high-performance liquid chromatography (HPLC) profile (s) to further confirm the active diterpenoid lactones present




Figure 2: Chemical structure of DDA. Molecular formula: $(C_{20}H_{28}O_4)$. Molecular weight: 332.44 g

AP chloroform extract (APCE), the following NMR analysis was carried out.

ANG is a bicyclic diterpenoid lactone and contains three hydroxyl groups at C-3, C-19, and C-14, these being secondary, primary, and allylic, respectively. It has been shown that the intact γ -butyrolactone ring, the double bonds at C-12 and C-13, C-8 and C-17, and C-14 hydroxyl group are responsible for the cytotoxic activity.^[13] The stereochemistry of ANG has previously been established. The two central six-membered rings adopt a chair conformation, and the furan ring adopts an envelope conformation.^[14-16] ANG forms hydrogen-bonded chains at both ends of the molecule.

DDA is another diterpenoid isolated from AP. It is a colorless needle crystal with the presence of hydroxyl, α , β -unsaturated- γ -lactone, and exomethylene groups in its chemical structure. This is very similar to that of DA, with the exception of a double bond at C-11 and C-12.^[17]


MATERIALS AND METHODS

1H -NMR spectra were recorded on BRUKER-400 spectrometer, NMR facility division, University Technology Mara, Shah Alam, Kuala Lumpur, Malaysia. Using $CDCl_3$ as a solvent and tetramethylsilane as an internal standard.

RESULTS AND DISCUSSION

The 1H NMR spectrum of chloroform fraction showed the presence of methyl group at H-18 (δ 1.289, 3H), H-20 (δ 0.801, 3H) and methylene group at H-11 (δ 2.353, 2H) and H-15 (δ 4.191, 2H). The presence of two double bonds was encountered at H-12 (δ 6.899, 1H) and H-17 (δ 4.598, 2H). The presence of a primary alcoholic group showed at H-19 (δ 3.352, 2H). These structural characteristic reveals the presence of ANG in APCE [Table 1.1].

The 1H NMR spectrum of chloroform fraction showed the presence of two methyl groups at H-18 (δ 1.684, 3H), H-20 (δ 0.801, 3H) and one methylene group at H-15 (δ 4.778, 2H). The three double bonds in the ring were ascertained at H-12 (δ 6.266, 1H), H-17 (δ 4.851, 2H), and H-14 (δ 7.271, 1H). The occurrence of primary alcoholic group is showed at H-19 (δ 4.533, 2H). The olefinic



APCE: *Andrographis paniculata* chloroform extract; ANG: Andrographolide

Proton	Chemical shift DDA		APCE		Standard DDA
	δ H	J (Hz)	δ H	J (Hz)	
3	3.510 (m, 1H)		3.795 (m, 1H)		3.76 m
11	6.90 (m, 1H)		7.172 (m, 1H)		7.23 m
12	6.141 (d, 1H)	12.0	6.266 (d, 1H)	16.0	6.20 d
14	7.271 (m, 1H)		7.271 (m, 1H)		7.27 m
15	4.795 (d, 2H)		4.778 (d, 2H)		4.77 d
17	4.824 (s, 2H)		4.851 (s, 2H)		4.85 s
18	1.568 (s, 3H)		1.684 (s, 3H)		1.62 s
19	4.540 (s, 2H)	12.0	4.533 (s, 2H), 3.3 3.79	12.0	4.52 s
20	0.829 (s, 3H)		0.801 (s, 3H)		0.94 s



linkage between H-11 and H-12 can be reconfirmed by the presence of single proton at H-11 (δ 7.172, 1H) and H-12 (δ 6.266, 1H). These structural interpretations assure the presence of DDA. The ^1H -NMR analysis reveals the significant presence of ANG and DDA in APCE [Table 1.2].

The aerial parts of the plant AP (leaves and stems) are used to extract the active phytochemicals using various solvents. Among all the fractions only chloroform fraction of AP was found to be more active in both *in vitro* and *in vivo* experiments. Previous investigations on the chemical composition of AP showed that it is a rich source of diterpenoids such as ANG, DA and DDA are pharmacologically active in treating various ailments, these compounds are present in a higher percentage in leaves [Figures 2-6]. From the NMR analysis, the highest percentage of the active diterpenoids including ANG, DA, and DDA were found in the chloroform extract of AP.

The study was supported by the Ministry of Science, Technology, and Environment for the research grant 304/PFARMASI/640043/k105 and 04/PFARMASI/6123003 for providing research chemicals and facilities.

Nil.

There are no conflicts of interest.

Figure 4: ¹H-NMR Spectral analysis of 14-deoxy-11, 12-didehydroandrographolideFigure 5: ¹H-NMR Spectral analysis of *Andrographis paniculata* chloroform extract




Figure 6: (a and b) ¹H-NMR Spectral analysis of *Andrographis paniculata* chloroform extract

REFERENCES

- Basak A, Cooper S, Roberge AG, Banik UK, Chrétien M, Seidah NG, *et al.* Inhibition of proprotein convertases-1, -7 and furin by diterpenes of *Andrographis paniculata* and their succinoyl esters. *Biochem J* 1999;338(Pt 1):107-13.
- Hidalgo MA, Romero A, Figueroa J, Cortés P, Concha II, Hancke JL, *et al.* Andrographolide interferes with binding of nuclear factor- κ B to DNA in HL-60-derived neutrophilic cells. *Br J Pharmacol* 2005;144:680-6.
- Zhang CY, Tan BK. Effects of 14-deoxyandrographolide and 14-deoxy-11,12-didehydroandrographolide on nitric oxide production in cultured human endothelial cells. *Phytother Res* 1999;13:157-9.
- Liu J, Wang ZT, Ji LL, Ge BX. Inhibitory effects of neoandrographolide on nitric oxide and prostaglandin E₂ production in LPS-stimulated murine macrophage. *Mol Cell Biochem* 2007;298:49-57. Polphana N, Rangkadilok N, Thongnest S, Ruchirawat S, Ruchirawat M, Satayavivad J, *et al.* Determination and variation of three active diterpenoids in *Andrographis paniculata* (Burm.f.) nees. *Phytochem Anal* 2004;15:365-71.
- Li W, Fitzloff JF. HPLC-PDA determination of bioactive diterpenoids from plant materials and commercial products of *Andrographis paniculata*. *J Liq Chromatogr Relat Technol* 2004;27:2407-20.
- Najib Nik A Rahman N, Furuta T, Kojima S, Takane K, Ali Mohd M. Antimalarial activity of extracts of Malaysian medicinal plants. *J Ethnopharmacol* 1999;64:249-54.
- Pramanick S, Banerjee S, Achari B, Das B, Sen AK Sr., Mukhopadhyay S, *et al.* Andropanolide and isoandrographolide, minor diterpenoids from *Andrographis paniculata*: Structure and X-ray crystallographic analysis. *J Nat Prod* 2006;69:403-5.
- Weiming C, Xiaotian L. Deoxyandrographolide-19 β -D-glucoside from the leaves of *Andrographis paniculata*. *Planta Med* 1982;45:245-6.
- Sharma A, Krishnan L, Handa SS. Standardization of the Indian crude drug Kalmegh by HPLC for the determination of andrographolide. *Phytochem Anal* 1992;3:129-31.
- Siripong P, Konkathip B, Preechanukool K, Picha P, Tunsuwan K, Taylor WC. Cytotoxic diterpenoid constituents from *Andrographis paniculata* nees leaves. *J Sci Soc Thailand* 1992;18:87-194.
- Cui L, Qiu F, Wang N, Yao X. Four new andrographolide metabolites in human urine. *Drug Metab Dispos* 2004;33:555-62.
- Nanduri S, Nyavanandi VK, Thunuguntla SS, Kasu S, Pallerla MK, Ram PS, *et al.* Synthesis and structure-activity relationships of andrographolide analogues as novel cytotoxic agents. *Bioorg Med Chem Lett* 2004;14:4711-7.
- Smith A, Toder BH, Carroll PJ, Donohue J. Andrographolide: An X-ray

- crystallographic analysis. J. Crystallogr Spectrosc Res 1982;12:309-19.
14. Fujita T, Fujitani R, Takeda Y, Takaishi Y, Yamada T, Kido M, *et al.* On the diterpenoids of *Andrographis paniculata* X-ray crystallographic analysis of andrographolide and structure determination of new minor diterpenoids. Chem Pharm Bull 1984;32:2117-25.
 15. Spek A, Duisenberg A, Labadie R, Ratnayake S, Abeysekera A, De Silva K. Hydrogen bonding andrographolide: 3-[2-[decahydro-6-hydroxy-5 (hydroxymethyl)-5,8a-dimethyl-2-methylene-1-naphthalenyl]-ethylidene] dihydro-4-hydroxy-2 (3H)-furanone. Acta Crystallogr 1987;c43:530-2.
 16. Zhang J, Yang C, Wu D. Study of chemical components in *Uncaria rhynchophylla* miq jacks. Chin Herb Med 1998;29:649-51.
 17. Akowagha GA, Zhari I, Norhyati I, Mariam A. HPLC and HPTLC densitometric determination of andrographolides and anti-oxidant potential of *Andrographis paniculata*. J Food Comp Anal 2006;19:118-26.