

Evaluation of Antioxidant and Analgesic Activities of Three Medicinal Plants

Amir Hossain, Akbar Hossain, Sultana Juhara Mannan

Department of Pharmacy, ASA University Bangladesh, Dhaka, Bangladesh

ABSTRACT


Background: Medicinal plant species with antioxidant and pro-oxidant properties contemplate in high number of flavonoids, carotenoids, and phenolic acids. This present study has been designed to evaluate antioxidant and analgesic activities of three Bangladeshi medicinal plants. **Materials and Methods:** In this study, after the phytochemical screening, antioxidant activity of the crude methanol extracts of three medicinal species naming *Bougainvillea glabra* (Nyctaginaceae), *Licuala grandis* (Arecaceae), and *Anthurium crystallinum* (Araceae) were evaluated using the spectrophotometric 1,1-diphenyl-2-picrylhydrazyl-free radical scavenging assay. In addition, total phenolic and total tannin contents have also been determined using standard gallic acid. Finally, analgesic activities of these plants have also been evaluated by acetic acid writhing inhibition and tail flick test in mice. **Results:** The inhibitory concentration 50% values of methanol crude extract of *B. glabra*, *L. grandis*, and *A. crystallinum* were 1.72, 2.18, and 13.21 $\mu\text{g/mL}$, whereas the standard (ascorbic acid) showed the value 3.90 $\mu\text{g/mL}$. The extracts of *B. glabra*, *L. grandis*, and *A. crystallinum* also showed the total phenolic contents as 40.97 ± 0.004 , 247.272 ± 0.016 , and 302.22 ± 0.011 mg gallic acid equivalent (GAE)/100 g, and the total tannin contents as 2172 ± 0.002 , 435.7 ± 0.01 , and 2414.98 ± 0.028 mg GAE/100 g of dried plant extract, respectively. The three-plant extracts have reduced the pain by both these methods in mice in a dose-dependent manner. **Conclusion:** In sum, it can be said that the three-plant species showed potent antioxidant and analgesic activities.

Key words: Analgesic, *Anthurium crystallinum*, antioxidant, *Bougainvillea glabra*, *Licuala grandis*

SUMMARY

- The *Licuala grandis* is a small palm tree, prospered in shaded location growing over 3 feet tall. On the other hand, *Anthurium crystallinum* is a flowering plant, native to rainforest margins and *Bougainvillea glabra* is an evergreen, climbing shrub with thorny stems that grows 9 m tall. These three plant species

have some traditional uses, for instance, wound healing, analgesic etc. In the present study, we confirmed some common phytochemicals such as flavonoids, tannins and alkaloids which indicate the presence of antioxidant and analgesic activities that resembles with our findings.



Abbreviations Used: DPPH: 1,1-diphenyl-2-picrylhydrazyl, ROS: Reactive oxygen species, RNS: Reactive nitrogen species, SEM: Standard Error of Mean, IC_{50} : Inhibitory concentration 50%, GAE: Gallic acid equivalent, 5-HT3: 5-hydroxytryptamine 3, 5-HT2A: 5-hydroxytryptamine 2A.

Correspondence:

Amir Hossain,
Lecturer, Department of Pharmacy, ASA
University, Dhaka, Bangladesh.
E-mail: amir09pharm@yahoo.com
DOI: 10.4103/pr.pr.164_18

Access this article online
Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

In living organisms, the both reactive oxygen species and reactive nitrogen species are known to cause damage to lipids, proteins, enzymes, and nucleic acids, leading to some disorders such as neurodegenerative disease, cancer, aging, malaria, atherosclerosis, diabetes, liver injury, Alzheimer, Parkinson, and some others pathological events.^[1,2] Finding antioxidant effects of different compounds derived from a number of plants drawing much more focus and interest that could be relevant in relation to their nutritional incidence and their role in health and disease.^[3-7] Several studies have shown the presence of different natural antioxidative compounds in various plant species. Largely, these compounds include phenolic- or nitrogen-containing compounds and carotenoids.^[8-10] Compounds with antioxidant activity act as pro-oxidants under certain conditions or in high concentrations. Prooxidant activity can accelerate oxidative damage to different cells.^[11-16] Potential antioxidant should therefore be tested for pro-oxidant activity as well.

In a study, *Bougainvillea glabra* has been shown to accumulate large amounts of flavanols.^[17] The antidiabetic and antilipidemic effects have also been evaluated for this plant species.^[18] It should be noted that no

report on antioxidant and analgesic activities were found. According to Hostettmann and Terreaux,^[19] the predictable number of higher plant species in the world is of about 400,000, by the same token, secondary metabolites from different plants are categorized by a vast chemical variety, and currently one-fourth of all prescribed pharmaceuticals complexes in developed countries are directly or indirectly (semi-synthetic) derived from these plants. As plants produce a huge number of antioxidants, they can represent a source of new compounds with antioxidant activities.^[20,21] Among the huge number of plant secondary metabolites, some compounds such as flavonoids indicate the presence of both antioxidant as well as analgesic activity.^[22] From this point of view, the

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Hossain A, Hossain A, Mannan SJ. Evaluation of antioxidant and analgesic activities of three medicinal plants. Phcog Res 2019;11:248-53.

main goal of this research was to study the antioxidant activities through 1,1-diphenyl-2-picrylhydrazyl (DPPH)-free radical scavenging method, and determination of total phenolic and tannin contents as well as analgesic activities of crude methanol extracts from three-plant species, namely *B. glabra*, *Licuala grandis*, and *Anthurium crystallinum*, belonging to the family Nyctaginaceae, Arecaceae, and Araceae, respectively.

MATERIALS AND METHODS

Collection of plant material

The extraction and other investigation processes were carried out on the bark of *B. glabra* and leaves of *L. grandis* and *A. crystallinum*. These plant samples were collected from botanical garden, Dhaka, Bangladesh, in September 2017. The collected plant parts were then carefully handled and any sort of undesirable materials were detached from the collected parts.

Drying and grinding

The collected plant samples were washed with water and allowed to dry for few weeks. When the plant parts were suitable for grinding, they were grinded into coarse powder. Finally, the powder material was stored in a sealed container and kept in a dark, cool, and dry place until farther processing.

Cold extraction

The finely powered three-plant parts were taken in separate clean glass containers and they were soaked with methanol. Then, the container was sealed and kept for 15 days. During this time, it was allowed to intermittent stirring and shaking. The mixture was then filtered by cotton. Finally, it was filtered by Whatman filter paper.

1,1-diphenyl-2-picrylhydrazyl-free radical scavenging assay

In vitro antioxidant activity of the three plant extracts was assessed by DPPH-free radical scavenging assay.^[23-26] To begin with, 0.004% DPPH solution (w/v) was prepared in ethanol and 3 mL of this solution was mixed with different concentrations of 1 mL ethanolic crude extract. Subsequently, the mixture was vigorously shaken and kept in a dark place for 30 min at room temperature. After that, absorbance was measured spectrophotometrically at 517 nm, and percentage inhibition was evaluated using following equation; % DPPH-free radical scavenging activity = $([A_0 - A_1]/A_0) \times 100$. Here, A_0 and A_1 indicate the absorbance of control or blank and absorbance of plant extract/positive control. The % scavenging activity was then plotted against log concentration and a graph for inhibitory concentration 50% (IC_{50}) was calculated.

Estimation of total phenolics

Total phenolic contents of all the extracts were assessed by Folin-Ciocalteu (FC) technique.^[27,28] At first, 2 mL FC (1/10) reagent was mixed with 2 mL sodium carbonate (75 g/L) and an aliquot of the extracts or positive control. The tubes were shaken for 15 s for the development of the color and then, the mixtures were allowed to stand 20 min at room temperature. Ultraviolet (UV) absorbance was then taken at 750 nm in a UV spectrophotometer. Then, a gallic acid standard calibration curve was developed having a positive control curve equation of $y = 0.062x - 0.0154$, $R_2 = 0.9818$ (*B. glabra*); $y = 0.0559x + 0.0728$, $R_2 = 0.9948$ (*L. grandis*); and $y = 0.072x - 0.0411$, $R_2 = 0.9734$ (*A. crystallinum*).

Determination of total tannins

Total tannin contents of the extracts were also determined FC reagent.^[29,30] To begin the process, 0.1 mL extracts and/or positive control was mixed

with 7.5 mL distilled water and 0.5 mL FC reagent. This preparation was then shaken vigorously and kept for 5 min. Then 35% sodium carbonate solution (1 mL) was added following the dilution with 10 mL distilled water. This preparation was then kept at room temperature for 30 min. UV absorbance was taken at 725 nm using UV spectrophotometer. Then, gallic acid standard calibration curve was developed and positive control curve equation was $y = 0.0219x - 0.0037$, $R_2 = 0.8626$ (*B. glabra*); $y = 0.0214x - 0.0034$, $R_2 = 0.8099$ (*L. grandis*); and $y = 0.0307x - 0.0159$, $R_2 = 0.9377$ (*A. crystallinum*).

Analgesic activity test

Acetic acid-induced writhing inhibition test

Analgesic activity test was evaluated using Swiss albino mice of both sexes (weighing about 25–40 g). All the animals were unfed for about 2 h before the start of the test. Mice were divided into four groups taking five mice in each group, where Group I has been received 1% Tween-80 solution in water (at dose of 10 mL/kg) and Group II has been received Diclofenac-Sodium (Na) as a positive control at a 25 mg/kg body weight dose. The other two groups, groups III and IV were treated with those three extracts at a dose of 250 and 500 mg/kg body weight, respectively. After 30 min, 0.7% acetic acid was injected intraperitoneally. Afterward, 5 min later writhing number was counted for 15 min.^[31]

Tail flick test

The effect of plant extracts on the latency of tail-flick response representing the pain threshold was investigated in mice with minor modification of the method of Yerima *et al.*^[32] The tail withdrawal from the heat (flicking response) was taken as the end point. A cutoff period of 10 s was kept to avoid damage to the tail. The drug was administered once daily for 5 consecutive days. On the 5th day, 1 h after drug administration, tail-flick response was recorded at 30, 60, 120, 180, and 240 min.

Statistical analysis

All analyses were double checked and passed in two replications. Mean $\pm$ standard error of the mean was used to the present data. All experimental parameters were evaluated for their significance level by correlation and regression analysis; the *t*-tests ($P < 0.05$) were used. Microsoft excel 2016 was used for both statistical analysis and graphical presentation.

RESULTS

Phytochemical screening

In phytochemical screening test, *B. glabra* showed the presence of reducing sugar, combined reducing sugar, tannins, flavonoids, and

Table 1: Result of phytochemical screening of methanol leaves extract of *Bougainvillea glabra*, *Licuala grandis*, and *Anthurium crystallinum*

Phytochemical group	<i>Bougainvillea glabra</i>	<i>Licuala grandis</i>	<i>Anthurium crystallinum</i>
Reducing sugar	+	+	+
Combined reducing sugar	+	+	-
Tannins	+	+	+
Flavonoids	+	-	+
Saponin	-	-	-
Gums	-	-	-
Steroids	-	+	+
Alkaloids	+	+	+
Glycoside	-	+	-
Proteins	-	+	+
Acidic compounds	-	+	+

+: Presence; -: Absence

alkaloids. On the other hand, *L. grandis* confirmed reducing sugar, combined reducing sugar, tannins, steroids, alkaloids, glycoside, proteins, and acidic compounds; and *A. crystallinum* revealed the presence of reducing sugar, tannins, flavonoids, steroids, alkaloids, proteins, and acidic compounds [Table 1].

1,1-diphenyl-2-picrylhydrazyl-free radical scavenging activity

Quantitative antioxidant activity was performed by DPPH-free radical scavenging assay where the IC_{50} values of methanol crude extract of *B. glabra*, *L. grandis*, and *A. crystallinum* were 1.72, 2.18, and 13.21 $\mu\text{g/mL}$, whereas the standard (ascorbic acid) showed the value as 3.90 $\mu\text{g/mL}$ [Figure 1].




Figure 1: Comparison of 1,1-diphenyl-2-picrylhydrazyl scavenging activity of *Bougainvillea glabra*, *Licuala grandis*, and *Anthurium crystallinum* with positive control (Ascorbic acid)

Total phenol content

Gallic acid equivalent (GAE) was used to determine and express total phenolic contents of the three extracts [Figure 2]. The total phenolic content of *B. glabra*, *L. grandis*, and *A. crystallinum* was 40.97 ± 0.004 , 247.272 ± 0.016 , and 302.22 ± 0.011 mg GAE/100 g of dried plant extract, respectively, as represented in Table 2.

Total tannin content

GAE was also used to determine the total tannin content of the three extracts [Figure 3]. Total phenolic contents of *B. glabra*, *L. grandis*, and *A. crystallinum* were 2172 ± 0.002 , 435.7 ± 0.01 , and 2414.98 ± 0.028 mg GAE/100 g of dried plant extract, respectively [Table 2].

Analgesic activity

Acetic acid-induced writhing inhibition test

The three extracts *B. glabra*, *L. grandis*, and *A. crystallinum* showed 60.68%, 54.71%, and 29.06% writhing inhibition at doses 500 mg/kg, and 38.46%, 38.46%, and 18.81% writhing inhibition at doses 250 mg/kg body weight, respectively. On the other hand, the standard Diclofenac-Na exhibited 70.01%, 70.09%, and 70.09% writhing reduction in mice [Table 3].

Tail flick test

In tail-flick response test, ibuprofen showed prolongation in the response time 1 h after drug administration. The three-plant extracts displayed analgesic activities in mice in a dose-dependent manner [Table 4]. However, it is important to say that they did not show any statistically significant differences.

DISCUSSION

Medicinal plants, certainly, possess a great potential to treat a number of mild-to-severe disorders. This is why, medicinal plants from all around




Figure 2: Determination of total phenolic contents of (a) *Bougainvillea glabra* (b) *Licuala grandis* and (c) *Anthurium crystallinum* with the help of gallic acid standard calibration curve

the world demonstrate a lot of pharmacological activities counting antioxidant, analgesic, anthelmintic, and antidiarrheal activities. All the three-plant extracts exhibited the presence of reducing sugars, tannins, and alkaloids. They also ensured the presence of some others secondary metabolites [Table 1]. On studying several literatures, it has been understood that the presence of flavonoids and tannins secondary metabolite in plants confirms the existence of antioxidants or free radical scavenging activity.^[33] Some other studies showed that

Table 2: Polyphenolic content of methanol leaves extract of *Bougainvillea glabra*, *Licuala grandis*, and *Anthurium crystallinum*

Polyphenolic compounds	Methanol crude extract	Content (mg GAE/100 g)
Total phenol	<i>B. glabra</i>	40.97±0.004
	<i>L. grandis</i>	247.272±0.016
	<i>A. crystallinum</i>	302.22±0.011
Total tannin	<i>B. glabra</i>	2172±0.002
	<i>L. grandis</i>	435.7±0.01
	<i>A. crystallinum</i>	2414.98±0.028

Each value represents the total content of the analysis±SEM. Expressed in terms of mg GAE/100 g dried plant extract. SEM: Standard error of mean; GAE: Gallic acid equivalent; *B. glabra*: *Bougainvillea glabra*; *L. grandis*: *Licuala grandis*; *A. crystallinum*: *Anthurium crystallinum*

flavonoids are responsible for analgesic^[34] and tannins for antidiarrheal and anthelmintic activities,^[34] whereas the presence of alkaloids in plants possess antibacterial activity.^[35]

The free radical scavenging activity of three-plant species growing in Bangladesh, *B. glabra*, *L. grandis*, and *A. crystallinum* belonging to the family *Nyctaginaceae*, *Arecaceae*, and *Araceae*, respectively, was determined consuming DPPH-free radical. The DPPH scavenger capacity of the extracts was compared with known antioxidative substances (ascorbic acid). The antioxidant activity is one of the most important pharmacological properties of plants. To test the free radical scavenging activity or the antioxidant activity of food and plant extracts, DPPH is widely used.^[36-38] which is readily scavenged by antioxidants.^[39] The scavenging ability of the extract was concentration dependent and expressed as IC₅₀ (Sample concentration required to decrease the initial concentration of DPPH by 50%) values. A higher antioxidant activity is indicated by a lower IC₅₀ value. In the present study, among the three-plant extracts, *B. glabra* and *L. grandis* showed a potent IC₅₀ values as 1.72 and 2.18 µg/mL compared to another plant extract, *A. crystallinum* showed 13.21 µg/mL, whereas the standard ascorbic acid showed the value of 3.90 µg/mL [Figure 1]. Comparing to the standard, the three crude extracts possess potent antioxidant activities. The result also suggests that the plant contains phytoconstituents which are capable of donating hydrogen to protect the cell from potential damage.

Table 3: Percentage analgesic activity of leaves extract of *Bougainvillea glabra*, *Licuala grandis*, and *Anthurium crystallinum* in acetic acid-induced pain model

Animal group	<i>Bougainvillea glabra</i>			<i>Licuala grandis</i>			<i>Anthurium crystallinum</i>		
	SE	Percentage writhing inhibition	t-test (P)	SE	Percentage writhing inhibition	t-test (P)	SE	Percentage writhing inhibition	t-test (P)
Negative control	3.14	0	-	3.141	0	-	3.141	0	-
Positive control (25 mg/kg)	1.04	70.1	4.9538 (0.0011)	1.049	70.09	4.9538 (0.0011)	1.048	70.09	4.95 (0.0011)
Extract (250 mg/kg)	1.86	38.46	2.466 (0.0390)	1.327	38.46	2.6402 (0.0297)	1.6	18.81	2.55 (0.034)
Extract (500 mg/kg)	0.58	60.68	4.4462 (0.0022)	0.509	54.71	4.0236 (0.0038)	1.536	29.06	1.94 (0.087)

Positive control used Diclofenac-Na. SE: Standard error




Figure 3: Determination of total tannin contents of (a) *Bougainvillea glabra* (b) *Licuala grandis* and (c) *Anthurium crystallinum* with the help of gallic acid standard calibration curve

Table 4: Analgesic activity of leaves extract of *Bougainvillea glabra*, *Licuala grandis*, and *Anthurium crystallinum* in tail flick test

Group (dose)	Duration of latency of tail-flick response (s) at various time intervals					
	Basal	30 min	1 h	2 h	3 h	4 h
Control (10 ml/kg)	4.19±0.31	4.01±0.30	4.27±0.34	4.45±0.52	4.35±0.74	4.28±0.37
Ibuprofen (100 mg/kg)	5.32±0.39	4.07±0.38	5.42±0.43	4.89±0.48	5.61±0.34	5.20±0.52
<i>B. glabra</i> (500 mg/kg)	4.08±0.39	4.45±0.28	4.22±0.38	3.83±0.31	4.39±0.30	4.47±0.28
<i>B. glabra</i> (250 mg/kg)	3.88±0.50	4.18±0.25	4.43±0.35	4.49±0.48	4.67±0.65	4.37±0.30
<i>L. grandis</i> (500 mg/kg)	4.36±0.51	4.24±0.39	4.51±0.44	4.76±0.33	5.19±0.23	3.56±0.60
<i>L. grandis</i> (250 mg/kg)	4.61±0.51	3.8±0.35	4.83±0.65	4.49±0.42	4.57±0.50	4.24±0.47
<i>A. crystallinum</i> (500 mg/kg)	4.34±0.42	3.74±0.50	4.75±0.32	4.23±0.79	4.34±0.36	4.83±0.44
<i>A. crystallinum</i> (250 mg/kg)	3.63±0.32	4.12±0.38	4.67±0.58	4.96±0.55	4.11±0.57	4.26±0.40

B. glabra: *Bougainvillea glabra*; *L. grandis*: *Licuala grandis*; *A. crystallinum*: *Anthurium crystallinum*

Plant phenolics such as flavonoids, phenolic acids, and tannins present in the fruits and vegetables have potential biological activities including anti-atherosclerotic, anti-cancer, and anti-inflammatory activities and such activities might be associated with their antioxidant activity.^[40] Plant phenolics containing hydroxyl groups have a good scavenging capacity.^[41,42] The extracts of *B. glabra*, *L. grandis*, and *A. crystallinum* also showed the total phenolic contents as 40.97 ± 0.004 , 247.272 ± 0.016 , and 302.22 ± 0.011 mg GAE/100 g of dried plant extract, respectively. The result proved that the extract possesses phenolic compounds which indicate the presence of hydroxyl groups [Table 2].

Bitter plant polyphenols such as tannins have astringent properties which mean that they can bind to protein molecules and precipitate or shrink them. Tannins contain sufficient hydroxyl groups and other free radicals for instance carboxyl and they bind proteins and other macromolecules. These free radicals protect cell damage.^[43] Some studies specify that tannins are effective against ulcerated or inflamed tissues and they also possess a good anticancer activity.^[44,45] The extract of the three plants, *B. glabra*, *L. grandis*, and *A. crystallinum* also showed the total tannin contents as 2172 ± 0.002 , 435.7 ± 0.01 , and 2414.98 ± 0.028 mg GAE/100 g of dried plant extract, respectively, which conclude that the plant extract contains a considerable amount of tannins which may be a source of free radicals that protect the cell from death and satisfy the possible mechanism how the plant extracts exerts its scavenging activities [Table 2].

In analgesic activity test, the intraperitoneal administration of acetic acid makes pain sensation^[46] and creates writhing reflex in animals by activating the chemosensitive nociceptors.^[47] Two different test methods have been employed in evaluating the analgesic activity of methanol extracts of three medicinal species. It is necessary to apply tests which differ with respect to stimulus quality, intensity, and duration, to obtain as complete a picture as possible of the analgesic properties of a substance using behavioral nociceptive tests.^[48] The results obtained indicate that the extract possesses a moderate dose-dependent analgesic effect on the various pain models used. Centrally acting analgesic drugs elevate the pain threshold of animals toward heat and pressure. The effect of the extract on these pain models indicates that it might be centrally acting.^[49] Acetic acid in this assay causes algesia by the release of some endogenous substances, which subsequently stimulated the pain nerve endings.^[50] It has also been noticed that in acetic acid-induced models, the analgesia level is reduced by a number of abdominal constrictions.^[51] It has been reported that flavonoids show analgesic action by increasing endogenous level of serotonin or interacting with 5-hydroxytryptamine 3 and 5-hydroxytryptamine 2A receptors.^[52] The three extracts showed significant writhing inhibition comparable to the standard drug Diclofenac-Na [Table 3]. The presence of flavonoid compounds in the plant extracts may be responsible for analgesic activity. The results showed a higher writhing inhibition at dose 500 mg/kg than dose 250 mg/kg body weight that was comparable to that of standard Diclofenac-Na. Among

the three extracts, *B. glabra* showed the highest writhing inhibition and all the results were statistically significant [Table 3].

CONCLUSION

The present study reveals that the methanol extracts of *B. glabra*, *L. grandis*, and *A. crystallinum* plants possess both antioxidant and analgesic activities. These preliminary studies do not describe the actual mechanism due to which the pharmacological activities were shown. However, a more extensive study (including the analysis and identification of the responsible compounds) is required to determine the exact mechanism of action of the extracts and their active compound(s) to authenticate them as a potent analgesic and antioxidant agent.

Acknowledgements

The authors are grateful to Department of Pharmacy, ASA University Bangladesh, for providing necessary instrumental support as well as chemicals and reagents.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Gutteridge JM. Biological origin of free radicals, and mechanisms of antioxidant protection. *Chem Biol Interact* 1994;91:133-40.
- Duan XJ, Zhang WW, Li XM, Wang BG. Evaluation of antioxidant property of extract and fractions obtained from a red alga, *Polysiphonia urceolata*. *Food Chem* 2006;95:37-43.
- Steinmetz KA, Potter JD. Vegetables, fruit, and cancer prevention: A review. *J Am Diet Assoc* 1996;96:1027-39.
- Aruoma OI. Free radicals, oxidative stress, and antioxidants in human health and disease. *J Am Oil Chem Soc* 1998;75:199-212.
- Bandoniene D, Pukalskas A, Venskutonis PR, Gruzdiene D. Preliminary screening of antioxidant activity of some plant extracts in rapeseed oil. *Food Res Int* 2000;33:785-91.
- Pieroni A, Janiak V, Dürr CM, Lüdeke S, Trachsel E, Heinrich M. *In vitro* antioxidant activity of non-cultivated vegetables of ethnic Albanians in Southern Italy. *Phytother Res* 2002;16:467-73.
- Couladis M, Tzakou O, Vervakidou E, Harvala C. Screening of some Greek aromatic plants for antioxidant activity. *Phytother Res* 2003;17:194-5.
- Shahidi F, Wanasundara PK. Phenolic antioxidants. *Crit Rev Food Sci Nutr* 1992;32:67-103.
- Velioglu YS, Mazza G, Gao YL, Oomah BD. Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *J Agric Food Chem* 1998;46:4113-7.
- Pietta P, Sionetti P, Mauri P. Antioxidant activity of selected medicinal plants. *J Agric Food Chem* 1998;46:4487-90.
- Cao G, Sofic E, Prior RL. Antioxidant and prooxidant behavior of flavonoids: Structure-activity relationships. *Free Radic Biol Med* 1997;22:749-60.
- Heijnen CG, Haenen GR, Oostveen RM, Stalpers EM, Bast A. Protection of

- flavonoids against lipid peroxidation: The structure activity relationship revisited. *Free Radic Res* 2002;36:575-81.
13. Amic D, Amic-Dragovic D, Bešlo D, Trinajstić N. Structure-radical scavenging relationship of flavonoids. *Croat Chem Acta* 2003;76:55-61.
 14. Subario A, Morita N. Prooxidant activity of lutein and its dimyristate esters in corn triacyl glyceride. *Food Chem* 2003;81:97-102.
 15. Young AJ, Lowe GM. Antioxidant and prooxidant properties of carotenoids. *Arch Biochem Biophys* 2001;385:20-7.
 16. Fukumoto LR, Mazza G. Assessing antioxidant and prooxidant activities of phenolic compounds. *J Agric Food Chem* 2000;48:3597-604.
 17. Heuer S, Richter S, Metzger JW, Wray V, Nimtz M, Strack D. Betacyanins from bracts of *Bougainvillea glabra*. *Phytochemistry* 1994;37:761-7.
 18. Adebayo GI, Alabi OT, Owoyele BV, Soladoye AO. Anti-diabetic properties of the aqueous leaf extract of *Bougainvillea glabra* (Glory of the Garden) on alloxan-induced diabetic rats. *Rec Nat Prod* 2009;3:187-92.
 19. Hostettmann K, Terreaux C. Search for new lead compounds from higher plants. *Chimia (Aarau)* 2000;54:652-7.
 20. Cuendet M, Potterat O, Salvi A, Testa B, Hostettmann K. A stilbene and dihydrochalcones with radical scavenging activities from *Loiseleuria procumbens*. *Phytochemistry* 2000;54:871-4.
 21. Bassman JH. Ecosystem consequences of enhanced solar ultraviolet radiation: Secondary plant metabolites as mediators of multiple trophic interactions in terrestrial plant communities. *Photochem Photobiol* 2004;79:382-98.
 22. Aiyegeoro OA, Okoh AI. Preliminary phytochemical screening and *in vitro* antioxidant activities of the aqueous extract of *Helichrysum longifolium* DC. *BMC Complement Altern Med* 2010;10:21.
 23. Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activities. *Lebensm Wiss Unterr Technol* 1995;28:25-30.
 24. Benabadi SH, Wen R, Zheng JB, Dong XC, Yuan SG. Anticarcinogenic and antioxidant activity of diindolylmethane derivatives. *Acta Pharmacol Sin* 2004;25:666-71.
 25. Sumi SA, Siraj MA, Hossain A, Mia MS, Afrin S, Rahman MM. Investigation of the key pharmacological activities of *Ficus racemosa* and analysis of its major bioactive polyphenols by HPLC-DAD. *Evid Based Complement Alternat Med* 2016;2016:3874516.
 26. Sánchez-Moreno C, Larrauri J, Saura-Calixto F. A procedure to measure the antiradical efficiency of polyphenols. *J Agric Food Chem* 1998;46:270-6.
 27. Boah H, Yadav RN, Unni BG. *In vitro* antioxidant and free radical scavenging activity of *Alternanthera sessilis*. *Int J Pharm Sci Rev Res* 2011;2:1502-6.
 28. Hossain A, Islam T, Saifuzzaman Saeed MA, Islam MK, Murshid GM, Rahman MM. Bioactivity of *Boehmeria macrophylla* (Urticaceae) leaf extract. *Orient Pharm Exp Med* 2016;16:233-41.
 29. Marinova D, Ribarova F, Atanasova M. Total phenolics and total flavonoids in bulgarian fruits and vegetables. *J Univ Chem Technol Metallurgy* 2005;40:255-60.
 30. Islam F, Hossain A, Hossain MG, Murshid GM, Rahman MM. Evaluation of antioxidant, analgesic and cytotoxic activities of *Typha angustata* L. Root. *Dhaka Univ J Pharm Sci* 2015;14:55-9.
 31. Ahmed F, Selim MS, Das AK, Choudhuri MS. Anti-inflammatory and antinociceptive activities of *Lippia nodiflora* Linn. *Pharmazie* 2004;59:329-30.
 32. Yerima M, Magaji MG, Yaro AH, Tanko Y, Mohammed MM. Analgesic and anti-inflammatory activities of the methanolic leaves extract of *Securinega virosa* (Euphorbiaceae). *Nig J Pharm Sci* 2009;8:47-53.
 33. Polterait O. Antioxidants and free-radical scavengers of natural origin. *Curr Org Chem* 1997;1:415-40.
 34. Mora A, Paya M, Rios JL, Alcaraz MJ. Structure activity relationships of polymethoxy flavones and other flavonoids as inhibitors of nonenzymic lipid peroxidation. *Biochem Pharmacol* 1990;36:317-22.
 35. Belemtougri RG, Constantin B, Cognard C, Raymond G, Sawadogo L. Effects of two medicinal plants *Psidium guajava* L. (Myrtaceae) and *Diospyros mespiliformis* L. (Ebenaceae) leaf extracts on rat skeletal muscle cells in primary culture. *J Zhejiang Univ Sci B* 2006;7:56-63.
 36. Cushnie TP, Cushnie B, Lamb AJ. Alkaloids: An overview of their antibacterial, antibiotic-enhancing and antivirulence activities. *Int J Antimicrob Agents* 2014;44:377-86.
 37. Da Porto C, Calligaris S, Celotti E, Nicoli MC. Antiradical properties of commercial cognacs assessed by the DPPH(.) test. *J Agric Food Chem* 2000;48:4241-5.
 38. Soares JR, Dinis TC, Cunha AP, Almeida LM. Antioxidant activities of some extracts of *Thymus zygis*. *Free Radic Res* 1997;26:469-78.
 39. Lu YR, Yeap Foo L. Antioxidant activities of polyphenols from sage (*Salvia officinalis*) *Food Chem* 2001;75:197-202.
 40. Chung KT, Wong TY, Wei CI, Huang YW, Lin Y. Tannins and human health: A review. *Crit Rev Food Sci Nutr* 1998;38:421-64.
 41. Kähkönen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS. Antioxidant activity of plant extracts containing phenolic compounds. *J Agric Food Chem* 1999;47:3954-62.
 42. Nacz M, Shahidi F. Extraction and analysis of phenolics in food. *J Chromatogr A* 2004;1054:95-111.
 43. Afify Ael-M, El-Beltagi HS, El-Salam SM, Omran AA. Biochemical changes in phenols, flavonoids, tannins, Vitamin E, β -carotene and antioxidant activity during soaking of three white sorghum varieties. *Asian Pac J Trop Biomed* 2012;2:203-9.
 44. Motar ML, Thomas G, Barbosa Fillo JM. Effects of *Anacardium occidentale* stem bark extract on *in vivo* inflammatory models. *J Ethnopharmacol* 1985;95:139-42.
 45. Ruch RJ, Cheng SJ, Klaunig JE. Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea. *Carcinogenesis* 1989;10:1003-8.
 46. Bauer AW, Kirby WM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 1966;45:493-6.
 47. Onasanwo SA, Elegbe RA. Antinociceptive and anti-inflammatory properties of the leaf extract of *Hedranthera barteri* in rats and mice. *Afr J Biomed Res* 2006;2:109-18.
 48. Tjølsen A, Berge OG, Hunskaar S, Rosland JH, Hole K. The formalin test: An evaluation of the method. *Pain* 1992;51:5-17.
 49. Adeyemi OO, Okpo SO, Okpaka O. The analgesic effect of the methanolic extract of *acanthus montanus*. *J Ethnopharmacol* 2004;90:45-8.
 50. Taesotikul T, Panthong A, Kanjanapothi D, Verpoorte R, Scheffer JJ. Anti-inflammatory, antipyretic and antinociceptive activities of *Tabernaemontana pandacqui* Poir. *J Ethnopharmacol* 2003;84:31-5.
 51. Marchiori M, Blank Mde F, Mourão RH, Antonioli AR. Anti-nociceptive activity of the aqueous extract of *Erythrina velutina* leaves. *Fitoterapia* 2005;76:637-42.
 52. Annegowda HV, Mordi MN, Ramanathan S, Mansor SM. Analgesic and antioxidant properties of ethanolic extract of *Terminalia catappa* L. Leaves. *Int J Pharmacol* 2010;6:910-5.