

## PHCOG RES.: Research Article

# Development and Evaluation of Polyherbal Formulations for Hair Growth Potential

Rathi Vaishali\*, Rathi Jagdish Chandra, Tamizharasi Sengodan<sup>1</sup>

*Department of Pharmacognosy, NRI Institute of Pharmaceutical Sciences, 3, Sajjan Singh Nagar, Bhopal - 462 026 India.*

*<sup>1</sup>Department of Pharmacognosy, Nandha College of Pharmacy, Perundurai Main Road, Erode - 638 052 India.*

\* Author for correspondence : E-Mail : roshan\_rathi2005@yahoo.co.in

### ABSTRACT

The present study is an effort to formulate and evaluate hair growth promoting activity of three polyherbal formulations. Polyherbal formulations were prepared using extract of *Cicer arietinum* Linn., *Ocimum sanctum* Linn. and *Cyperus rotundus* Linn. in various ratios to obtain the best formulation. The extract incorporated into cream were applied topically on shaved skin of rats and primary skin irritation test, hair growth initiation time, completion time, hair length and diameter were recorded. The ratio of *Cicer arietinum* Linn., *Ocimum sanctum* Linn. and *Cyperus rotundus* Linn. in 1:2:3 showed excellent hair growth activity comparable to standard.

**Keywords:** Alopecia, Polyherbal, Hair growth, Herbal formulation.

### INTRODUCTION

Alopecia is a universal problem, having affected both sexes of all races to different extents for as long as mankind has existed. It has been suggested that alopecia could have an adverse effect on physiological life and self esteem between both the genders (1). Alopecia affects approximately 50% of men over 40 years of age and may also affect just as many as women. The majority of men and women (90%) or more want to reverse, halt hair loss. Alopecia is a synonym of baldness, involves absence or loss of hair, especially of the head. Androgens are well known to cause regression and balding on the scalp in genetically disposed individuals. Alopecia has also been observed as major side effect of anticancer drugs, immunosuppressant and many other drug treatments. Minoxidil, a drug of scientific origin was scientifically proved for the treatment of alopecia (2). Though the

side effect associated with this drug has limited its pharmacological benefits hence the drug of plant origin is necessary to replace the synthetic one. India is a repository of medicinal plants (3–4). Besides healthcare, herbs are also used for beautification of the body and for preparation of various cosmetics (5). In traditional system of medicine, many plants and herbal formulations are reported for hair growth promotion (6–10) but lack of sound scientific backing and information limits their use.

The present study is an effort to formulate and evaluate hair growth promoting activity of polyherbal formulation, which include various concentrations of *Cicer arietinum* Linn., *Ocimum sanctum* Linn. and *Cyperus rotundus* Linn. The herbs *Cicer arietinum* Linn. and *Cyperus rotundus* Linn. were selected on the basis of their traditional use (11–12) and *Ocimum sanctum* Linn. was selected based on its anti-androgenic property (13).

## MATERIALS AND METHODS

### Plant material

The leaves of *Cicer arietinum* Linn., whole plant of *Ocimum sanctum* Linn. and roots of *Cyperus rotundus* Linn. were collected in the month of October locally from Bhopal. The plants were authenticated at Department of Pharmacy, Barkatullah University, Bhopal. The plants were dried under shade.

### Extraction

Dried powdered drug were taken and maceration was done by keeping them in 95% alcohol for 7 days with occasional stirring. After filtration, double maceration was done for next three days with 95% alcohol. The solvent was removed under reduced pressure and the extract obtained was air dried.

### Formulation

Herbal hair creams were prepared by fusion method using o/w type base (14). The formula of base contains glyceryl mono stearate 9% w/w, liquid paraffin (light) 20% w/w, cetyl alcohol 15% w/w, beeswax 15% w/w, propyl and methyl paraben 0.15% w/w, glycerol 4.5% w/w and water 59% w/w. The 5% extract mixture of *Cicer arietinum* Linn., *Ocimum sanctum* Linn. and *Cyperus rotundus* Linn. in various ratio as shown in Table 1 were incorporated in the base to obtain 5% herbal cream F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> respectively.

### Animals

Albino rats, weighing 120- 150 g were used for hair growth activity. The study was approved by Institutional Animal Ethical Committee, Barkatullaha University, Bhopal. Animals were placed in cages and kept in standard environmental conditions, fed with standard diet ad libitum and allowed free access to drinking water. The prepared formulations were assessed for hair growth studies.

### Primary skin irritation test

The rats were divided into five groups of six rats each. A 4cm<sup>2</sup> area of dorsal portion of all the rats were shaved and wiped with surgical spirit. Measured quantity of

formulation of F<sub>1</sub>, F<sub>2</sub>, and F<sub>3</sub> were applied over the site. The test sites were observed for erythema and edema for 48 h after application (15).

### Hair growth activity

The rats were divided into five groups of six rats each. A 4 cm<sup>2</sup> area of dorsal portion of all the rats were shaved and wiped with surgical spirit. Hair remover was also applied over the shaved area to assure the removal of trace of hairs from denuded area. Group1 was kept as control, where there was no drug treatment. Group2 was treated as standard, where 2% minoxidil lotion (Mintop) was applied over the shaved area, once a day. The animals of remaining groups were given application of 5% cream of formulation, F<sub>1</sub>, F<sub>2</sub> and F<sub>3</sub> respectively. The treatment was continued for 30 days during which time, hair growth initiation (minimum time to initiate hair growth on denuded skin region) and completion time (time taken to completely cover the denuded skin region with new hair) were recorded for each group of animals (16). Hair was plucked randomly from the shaved area of selected rats, from each group on 10<sup>th</sup>, 20<sup>th</sup> and 30<sup>th</sup> day of the treatment and length and diameter of 24 hairs was measured (17). The average length and diameter were determined. The determination and evaluation of these parameters have been considered as vital for accomplishment of hair growth. It is considered that these parameters are accomplishing the concept of hair growth.

## RESULTS AND DISCUSSION

### Primary skin irritation test

This test was conducted to evaluate the irritancy of the prepared formulations on intact skin of rats. None of the prepared formulations showed any erythema or edema, indicating that the prepared formulations were non-irritant on the skin of rats.

### Hair growth activity

In control group animals, initiation of hair growth in denuded area was observed in second week. Hair growth initiation was noted in the first week in rats of minoxidil treated standard group. The formulation F<sub>3</sub> exhibited hair growth initiation on 7<sup>th</sup> day and F<sub>2</sub> on 9<sup>th</sup> day whereas with formulation F<sub>1</sub>, hair growth initiation time was reduced to 4<sup>th</sup> day. Similarly the time taken for complete hair growth on shaved area was affected with minoxidil treatment as well as treatment with formulations. Complete hair growth with minoxidil and control group was observed in 20 and 24 days respectively. In formulation F<sub>2</sub>

**Table 1. Various ratios of extract for formulations**

Formulation	<i>Cicer arietinum</i> Linn	<i>Ocimum sanctum</i> Linn	<i>Cyperus rotundus</i> Linn
F1	1	2	3
F2	2	3	1
F3	3	1	2

**Table 2. Effect of various formulations on hair growth initiation and completion time of albino rats**

Compound Name	Initiation Time (in days)	Completion Time (in days)
Control	10.67 ± 1.37	24.83 ± 2.14
Standard	6.17 ± 1.47**	20.00 ± 2.61**
F <sub>1</sub>	4.17 ± 1.17**	18.17 ± 1.47**
F <sub>2</sub>	9.17 ± 1.47	22.83 ± 1.60
F <sub>3</sub>	6.83 ± 1.47**	19.83 ± 1.72**

Value are mean ± S.D. ;

\*P<0.01,

\*\*P<0.001, When compare to control value by student's t-test (n = 6 animals)

**Table 3. Effect of various formulations on hair length of albino rats**

Compound Name	Length of hair in mm		
	10 Days	20 Days	30 Days
Control	3.00 ± 0.72	6.17 ± 0.76	7.54 ± 0.83
Standard	3.46 ± 0.59 *	6.79 ± 0.98 *	8.5 ± 1.10 **
F <sub>1</sub>	4.08 ± 0.72 ***	7.33 ± 1.2**	9.08 ± 1.14***
F <sub>2</sub>	3.29 ± 0.69	6.46 ± 1.06	7.88 ± 1.15
F <sub>3</sub>	3.67 ± 0.70**	6.96 ± 1.23*	8.38 ± 1.28 **

Value are mean ± S.D.

\*P<0.05;

\*\*P<0.01;

\*\*\*P<0.001; When compare to control value by student's t-test (n = 24 hairs)

**Table 4. Effect of various formulations on hair length of albino rats**

Compound Name	Diameter of hair in mm		
	10 Days	20 Days	30 Days
Control	0.0192±0.0065	0.0233±0.007	0.0279±0.0066
Standard	0.0246±0.0083*	0.0358±0.0102 ***	0.0458±0.0102***
F <sub>1</sub>	0.0271±0.0127**	0.0383±0.0096***	0.0479±0.0093***
F <sub>2</sub>	0.0208±0.0097	0.0254±0.0078	0.0304±0.0069
F <sub>3</sub>	0.0254±0.0078**	0.0350± 0.006***	0.0408±0.0102 ***

Value are mean ± S.D.

\*P<0.05;

\*\*P<0.01;

\*\*\*P<0.001; When compare to control value by student's t-test (n = 24 hairs)

complete hair growth occurred after 22 days, in F<sub>3</sub> after 19 days, and in F<sub>1</sub> it was reduced to 18 days (Table 2). In comparison to control for formulation F<sub>1</sub> the whole denuded area was covered with hair during the 4<sup>th</sup> week (Fig. 1). The experiment thus clearly demonstrates hair growth promoting activity in the developed formulations. The length of the hair began to increase until the end of the treatment course (Table 3). The formulation F<sub>1</sub> produced a greater effect on the length of hair when compare to other group being 9.08 mm at the end of the course, compare to 7.88 mm in the F<sub>2</sub>, 8.38 mm in F<sub>3</sub> and 8.5 mm in standard. This may be due to the premature



**Figure 1:** (A) Initially shaved albino rat (B) Control group after 30 days (C) Albino rat treated with formulation F1 for 30 days showing complete hair growth.

switching of follicles from the telogen to anagen phase of hair growth cycle (18). On 30<sup>th</sup> day, in control animal diameter of hair was found 0.0279 mm, in standard it was 0.0458 mm, in formulation F<sub>2</sub> and F<sub>3</sub> it was 0.0304 mm and 0.0408 mm respectively, but in formulation F<sub>1</sub>, it was highest around 0.0479 mm (Table 4).

## CONCLUSION

Among the various formulations, the formulation F<sub>1</sub>, showed better growth initiation and hair growth completion time at the same time formulation F<sub>1</sub>, showed remarkable improvement in hair length and diameter compare to control, standard and other formulations. Hence it can be concluded that the extract combination F<sub>1</sub> proved excellent growth activity and the formulation F<sub>1</sub>, might be hold a promise of potential herbal alternative for synthetic drugs used for alopecia.

## REFERENCES

1. Bhalerao S.S. and Solanki N.H. Therapeutic approaches to management of common baldness. *Indian Drug*. **39**(11): 567–573 (2002).
2. Reddy M.S., Mautalik S. and Rao V.G. Preparation and evaluation of minoxidil gels for topical application in alopecia. *Indian J Pharm Sci*. **68**(4): 432–436 (2006).
3. Saraf S., Pathak A.K. and Dixit V.K. Hair growth promoting activity of *Tridax procumbens*. *Fitoterapia*. **62**: 495–498 (1991).
4. Ansari S.H. and Ali M. Hair care and herbal drug. *Indian J Nat Prod*. **13**(1): 3–5 (1997).
5. Aburjai T. and Natsheh F.M. Plants used in cosmetics. *Phytother Res*. **17**: 987–1000 (2003).
6. Matsuda H., Yamazaki M., Asanuma Y. and Kubo M. Promotion of hair growth by *Ginseng radix* on cultured mouse vibrissal hair follicles. *Phytother Res*. **17**: 797–800 (2003).
7. Grindlay D. and Reynolds T. The aloe vera phenomenon: a review of the properties and modern uses of the leaf parenchyma gel. *J Ethnopharm*. **16**: 117–151 (1986).
8. Dweck A.C. African plants. *Cosmet Toiletries*. **112**: 41–51 (1997).
9. Kamimura A. and Takahashi T. Procyanidin B-2, extracted from apples, promote hair growth: a laboratory study. *British Journal of Dermatology*. **146**: 41–51 (2002).
10. Rathi V., Rathi J.C., Tamizharasi S. and Pathak A.K. Plants used for hair growth promotion: A review. *Phcog Rev*. **2**(3): 165–167 (2008).
11. Kirtikar K.P. and Basu B.D., Indian Medicinal Plants, Vol I, (International Book Distributors, Dehradun, 1995) 768–769.
12. Ayurvedic Pharmacopocia of India, Vol III, (Govt of India, Ministry of Health and Welfare, New Delhi, 1999) 129.
13. Das S.K. and Vasudevan D.M. Tulsi: The Indian holy power plant. *Nat Prod Rad*. **5**(4): 279–280 (2006).
14. Dixit V.K., Adhirajan N. and Gowri C. Development and evaluation of herbal formulations for hair growth. *Indian Drugs*. **38**(11): 559–563 (2001).
15. Patni P., Varghese D., Balekar N. and Jain D.K. Formulation and evaluation of herbal hair oil for alopecia management. *Planta Indica*. **2**(3): 27–30 (2006).
16. Roy R.K., Thakur M. and Dixit V.K. Effect of *Cuscuta reflexa* Roxb. On hair growth activity of albino rats. *Indian Drugs*. **43**: 951–956 (2006).
17. Adirajan N., Ravikumar T., Shanmugasundaram N. and Babu M. *In vivo* and *in vitro* evaluation of hair growth potential of *Hibiscus rosa sinensis* Linn. *J Ethnopharm*. **88**: 235–239 (2003).
18. Philpot M.P., Green M.R. and Kealey T. Rat hair follicle growth in vitro. *Br J Dermatol*. **127**: 600–607 (1992).