

Inter Simple Sequence Repeat Analysis of Genetic Diversity and Relationship in Four Egyptian Flaxseed Genotypes

Abeer A. El Sayed¹, Shahira M. Ezzat^{1,2}, Saber H. Mostafa³, Saeid Z. Zedan³, Essam Abdel-Sattar¹, Nebal El Tanbouly¹

¹Department of Pharmacognosy, College of Pharmacy, Cairo University, Cairo 11562, ²Department of Pharmacognosy, Faculty of Pharmacy, October University for Modern Science and Arts, 6th October 12566, ³Crops Research Department, Agricultural Research Center, Giza, Egypt

ABSTRACT

Background: Flaxseed is a highly important industrial and medicinal plant worldwide. **Objective:** To use inter simple sequence repeat (ISSR) technique for making unique fingerprint for the four newly produced genotypes of flax in Egypt. **Materials and Methods:** The genetic diversity among four promising Egyptian flax (*Linum usitatissimum* L.) genotypes was premeditated by means of polymerase chain reaction-based ISSR markers. The phenotypic variation among the four flax genotypes, namely, promising strains 533/39/5/3 (F1), S.402/3/3/7 (F2), S.421/3/6/4 (F3), and S.11 (F4) was studied during the two successive winter seasons of 2014/2015 and 2015/2016 in randomized complete block design through four replications. **Results:** The promising strain (F3) surpassed the other flax genotypes regarding seed yield/feddan, oil yield/feddan, and oil percentage. Twelve ISSR primers were used for the genetic examination yielding 139 loci, of which 31 were polymorphic. The middling number of amplified loci and the middling number of polymorphic loci per primer were 11.6 and 2.6, correspondingly, while the percent of loci polymorphism ranged from 0.0% to 58.0% with a middling of 21.4% crosswise all the flax genotypes. The more informative primers were GAC (GATA)₄ and (GATA)₄ GC, while the less informative were (AC)₈T and (GT)₈G. Unweighted pair group method with arithmetic mean derived dendrogram clearly discriminated the flax genotypes in three clusters. The Jaccard's similarity coefficient along with the genotypes ranged from 0.91 to 0.95. **Conclusion:** This study identified S. 421/3/6/4 (F3) strain to be the mainly assorted genotype and recommended its use in propagation programs and for upward mapping populations.

Key words: Cluster analysis, genotypes, linseed, molecular marker, polymorphism


SUMMARY

Four promising Egyptian flax (*Linum usitatissimum* L.) genotypes was

studied using ISSR. The more informative primers were GAC(GATA)₄ and (GATA)₄ GC. The strain S. 421/3/6/4 (F3) was identified as the most diverse genotype.



Linum usitatissimum L.



Clustering of 4 Flax cultivars

Abbreviations Used: CTAB: N-cetyl-N,N,N-trimethylammonium bromide; EDTA: ethylenediaminetetraacetic acid; ISSR: Inter Simple Sequence Repeat; PCR: polymerase chain reaction; RAPDs: random amplified polymorphic DNAs; RCBD: Randomized Complete Block Design; UPGMA: Unweighted Pair Group Method with Arithmetic Mean; SSR: simple sequence repeat.

Correspondence:

Dr. Shahira M. Ezzat,
Department of Pharmacognosy,
College of Pharmacy, Cairo University,
Cairo P.O. Box 11562, Egypt.
Department of Pharmacognosy, Faculty of
Pharmacy, October University for Modern Science
and Arts, 6th October, P.O.Box 12566, Egypt.
E-mail: shahira.ezzat@pharma.cu.edu.eg
DOI: 10.4103/pr.pr_126_17

Access this article online

Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

Family Linaceae encompass 22 genera, of which genus *Linum* is the majority well known. This genus includes the refined species *Linum usitatissimum* and the ornamentals *Linum grandiflorum* and *Linum perenne*. However, the concluding two species are of little economic importance.^[1]

Flax (*L. usitatissimum* subsp. *usitatissimum*) is one of the beginning crops in Egypt cultivation is a diploid, annual plant variety, which is mainly measured to be inbreeding. Ever since the domestication of flax, there has been a partiality for mounting it either for its fiber or oil. The twofold idea of flax was already known in ancient Egypt, linen (consequent from the fiber) was used for covering the regal mummies, and as well linseed oil was old to embalm the bodies of late Pharaohs.^[2]

Over the last two decades, there has been rehabilitated attention in the use of flaxseed for its fiber, oil, and functional food production.^[3] Being a highly industrial and medicinal plant, the worldwide demand of flaxseed species is increasing day by day; therefore, more researches have been conducted in the development of linseed genotypes. Miscellany

assessment of flax was former attempted by means of morphological parameters^[4] and isozyme markers.^[5]

Molecular categorization of flax germplasm has been finished through various molecular techniques to assess genetic diversity of the cultured flax and to inspect evolutionary dealings of undomesticated flax species.^[6] The use of DNA-based markers to learn flax diversity was early reported by Oh *et al.* (2000).^[4] Among the molecular markers, microsatellites called simple sequence repeat (SSR) markers were the most suitable for several applications because of the effortlessness in

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: El Sayed AA, Ezzat SM, Mostafa SH, Zedan SZ, Abdel-Sattar E, El Tanbouly N. Inter simple sequence repeat analysis of genetic diversity and relationship in four egyptian flaxseed genotypes. Phcog Res 2018;10:166-72.

handling, reproducibility, co-dominant inheritance, and genome-wide coverage.^[7] Inter SSR (ISSR) region can be amplified using different protocols. Protocol of ISSR-polymerase chain reaction (PCR) for agarose gel electrophoresis is a fast and efficient technique for standardizing primers and quick polymorphism testing including mapping. Microsatellites are highly useful markers in cultivars identification as they have been shown to be highly polymorphic and genotype specific. Thus, ISSR technique is highly robust and reproducible; hence, it was rapidly replacing random amplified polymorphic DNAs (RAPDs) as a tool for genotype segregation.^[8]

Other studies were carried out on genetic assortment of flax varieties through RAPD technique. DNA (RAPD) markers were used to cram multiplicity in 61 flax varieties including Canadian cultivars and landraces, where low genetic unpredictability was reported.^[9] RAPD markers were also used to analyze the genetic difference, genetic attrition, and connection in 54 North American flax cultivars.^[10] Furthermore, phenotypic and RAPD disparity inside four infraspecific groups of flax (dehiscent flax, fiber flax, large-seeded flax, and intermediate flax) was studied to recognize phenotypic and genotypic demarcation within the cultivated gene pool.^[11]

The ISSR practice for flax fingerprinting was optimized using re-amplification technique and through arithmetical correlation of free energy of dissociation of ISSR primers.^[12,13] An ISSR primer assay was reported in the cram of flax germplasm.^[14,15] The molecular genetic analysis realized by the ISSR method made it possible to obtain the objective data on genetic relations between flax genotypes.^[15] Fingerprinting the marketable flax genotypes based on molecular markers is a crucial measure for unambiguous and quick identification of similar or closely related genotypes.

Flaxseed oil is becoming increasingly popular as a nutritional and functional food in the Western world due to its high content of therapeutic health-promoting substances such as omega-3 fatty acid, soluble and insoluble fiber, and lignans, and its suitability for use in bread, breakfast cereals, muesli bars, and other food products.^[16] Flaxseed straw contains bast fibers which can be used for the production of paper, coarse textiles, rope, fiber board, molded panels, and as insulation material.^[17,18] Hence, there has been increased interest in 1 breeding and growing dual-purpose linseed cultivars which can be harvested for both seed and fiber.^[19]

In the current study, we planned to investigate the difference between the commercial genotype Sakha 1 and the four promising flax genotypes concerning fiber and oil yields. In addition, the ISSR technique was chosen for making unique fingerprint for the four newly produced genotypes of flax to evaluate the genetic diversity among them, with the aim of providing facts and tools to increase the assortment for future flax propagation and assisting in developing and planning breeding strategies for crop improvement programs in Egypt.

MATERIALS AND METHODS

Plant germplasm and phenotyping

Two field experiments were conducted at Giza Agriculture Research Station, Giza Governorate, Egypt, during 2014/2015 and 2015/2016 winter seasons to study the performance of five flax genotypes concerning straw, fiber seed, and oil yields as well as their related characters. The genotypes included the local cultivar Sakha 1 as well as the promising strains 533/39/5/3 (F1), S.402/3/3/7 (F2), S.421/3/6/4 (F3), and S. 11 (F4) [Table 1]. The experiments trails were arranged in a randomized complete block design with four replications. Sowing date was in the first week of November in both seasons; the plot size was 10.5 m². Plant density of 2500 seeds/m² was used and seeds were broadcasted regularly within each plot. Normal cultural practices for flax production as

Table 1: Pedigree, classification, and characteristic of four Egyptian flax genotypes

Genotype	Type	Pedigree
Sakha 1	Dual	Bombay X I.1485
533/39/5/3 (F1)	Fiber	S.420/140/5/10 X Bombay
S.402/3/3/7 (F2)	Dual	235 X Giza 5
S.421/3/6/4 (F3)	Dual	S.162/12 X S. 6/2
S.11 (F4)	Fiber	H. (420/140/5/10 X S, 401/2) X I, 1563

recommended were followed.^[20] At maturity, straw, seed, fiber, and oil yields/fad were calculated from the hole plot area basis. The seeds of the four promising flax genotypes are held in reserve in the herbarium of the Pharmacognosy Department, College of Pharmacy, Cairo University as Voucher specimen no. 2092016. Samples of the leaves, obtained from seedlings of the germinated seeds, were stored at -70°C, freeze-dried, and ground to a well grind using a coffee chopper before DNA isolation. Studied characters were straw yield/feddan (t.); seed yield/feddan (kg); fiber yield/feddan (kg); oil yield feddan; fiber percentage (%); and oil percentage (%).

Materials for DNA mapping

Buffers: The following buffers were used: Extraction buffer: 0.7 M NaCl, 100 mM Tris (pH 7.5), 0.01 M ethylenediaminetetraacetic acid (EDTA), 1% (w/v) N-cetyl-N, N, N-trimethylammonium bromide (CTAB), and 1% (v/v) β-mercaptoethanol (added immediately before use); washing buffer: 1:76% ethanol, 0.2 M Na-acetate; 2:76% ethanol as washing buffer, 10 mM NH₄ O-acetate, TE-buffer; 10 mM tris (pH 8.0), 1 mM EDTA, ×10; reaction buffer: 100 mM tris (pH 8.3), 500 mM KCl, 0.01% (w/v) gelatin, chloroform/isoamyl alcohol 24:1 (v/v), isopropanol, d NTP, Taq DNA polymerase.

Primers: Twelve primers, ISSR (Applied Biosciences), were used in the detection of polymorphism and used in the present analysis [Table S1]. ISSR practice was carried out in triplicates using genomic DNA with 12 decamer primers for reproducibility of the consequences.

Molecular weight markers: 100 bp ladder (Promega Corporation, USA).

Equipment

ADNA thermocycler (Hybaid PCR Express, USA) was used for the amplification of DNA, and agarose gel electrophoresis implement (Biorad Wide Mini Sub Cell, USA) was used for the separation of ISSR fragments according to size and ultraviolet (UV) Polaroid camera used for the apparition of ISSR wreckage.

Methods for molecular investigations

DNA extraction

DNA analysis was conducted at Food Technology Research Institute, Agriculture Research Center, Ministry of Agriculture and Land Reclamation, Giza, Egypt, in 2016. DNA was extracted using CTAB method.^[21] Fifty milligrams of frozen leaf were pulverized in liquid nitrogen, extracted with 0.8 ml CTAB, and precipitated with isopropanol.

Assessment of DNA deliberation

DNA concentration was determined by diluting the DNA 1:5 in distilled H₂O. The DNA samples were electrophoresed in 1% agarose gel against 10 µg of a DNA size marker. This marker covers a range of concentration between 95 ng and 11 ng. Thus, valuation of the DNA concentration in a prearranged sample was achieved by comparing the intensity of fluorescence of the unknown DNA band with the dissimilar bands in the DNA size marker.

Magnification of inter simple sequence repeat markers

The PCRs were conceded out using 100 ng of genomic DNA template subsequent a thermal cyclic program.^[22]

Thermocycling profile

Magnification of PCR was performed in a Perkin-Elmer/GeneAmp® PCR System 9700 (PE Applied Biosystems, USA) automatic to accomplish 35 cycles later than an early denaturation cycle for 5 min at 94°C. Each cycle consisted of a denaturation step at 94°C for 1 min, an annealing step at 45°C for 1 min, and an elongation step at 72°C for 1.5 min. The primer expansion segment was wholesale to 7 min at 72°C in the closing cycle. The augmentation products were determined by electrophoresis in a 1.5% agarose gel containing ethidium bromide (0.5ug/ml) in 1 × TBE buffer at 95 volts. A 1 kb DNA ladder was used as a molecular size standard. PCR products were visualized on UV light and photographed using a Gel Documentation System (Bio-Rad 2000, Germany).

Inter simple sequence repeat – polymerase chain reaction analysis

DNA was extracted from fresh leaves using the QiagenDNeasy kit. PCR magnification was performed in a Perkin-Elmer/GeneAmp® PCR system 9700 (PE Applied Biosystems, USA). Intensification products were determined by electrophoresis, visually examined and scored for the occurrence (1) or absence (0) of DNA bands. The resemblance matrix was obtained through the cluster analysis of data using unweighted pair-group method with arithmetic average (UPGMA). For estimating genetic distance among the tested samples, each of DNA bands was treated as a unit character.^[23] A dendrogram was constructed using the UPGMA with the SAHN module of NTSYS-pc to show a phenetic representation of genetic relationships as exposed by the similarity coefficient.^[24]

Data analysis

Every data were statistically analyzed by the study of variance method according to Snedecor and Cochran.^[25] Differences between means were tested by least significant difference at the level of 0.05. Bartlett test of homogeneity was adapted indicating no statistical evidence for heterogeneity, thus combined analysis of variance (ANOVA) for genotypes over seasons was worked out according to Le Clerg *et al.*^[26]

The banding patterns generated by ISSR-PCR marker analyses were scored as present (1) or absent (0), each of which was treated as a sovereign character in spite of its intensity. Only major and reproducible bands obtained for each ISSR primer were measured. By comparing the banding patterns of species for a primer, species-specific bands were identified. Faint or indistinct bands were not measured.

RESULTS

Phenotypic variations, correlations, and analysis of variance for the four genotypes

Statistical analysis of all phenotypic data was conducted to account for the phenotypic variation characters of the four promising Egyptian flax genotypes [Table 1]. Large phenotypic variations were observed concerning the yields (mean values) of straw, seed, fiber, and oil/feddan (fed.) as well as fiber and oil percentages. Results obtained from the combined analysis over the two winter seasons 2014–2015 and 2015–2016 are presented in [Table 2]. Data offered in Table 2 revealed significant differences among the four flax genotypes regarding all the six characters (straw yield/fed., seed yield/fed., fiber yield/fed., oil yield/fed., and fiber and oil percentage). ANOVA indicated that the promising strain S. 533/39/5/3 (F1) ranked first in the yields of straw, fiber, and fiber percentage with values of 4.70 T/fed., 814.35 kg/fed., and 18.30%, respectively. The superiority ratios of S. 533/39/5/3 (F1) over Sakha 1 were 28.77%, 36.40%, and 11.86% for the previously mentioned characters, respectively.

On the other hand, the promising strain 421/3/6/4 (F3) gave the maximum estimates for seed yield (743.00 kg/fed.), oil (304.85 kg/fed.), and oil percentage (41.03%); the superiority ratios of S. 421/3/6/4 (F3) over the variety Sakha 1 were 17.94%, 21.89%, and 3.35% for seed kg/fed., oil kg/fed., and oil percentage, respectively. The other flax genotypes laid intermediate position between highest estimates and the lowest one.

Polymorphism revealed by inter simple sequence repeat analysis

The banding profile of the four promising genotypes of flax produced by the 12random primers is illustrated in Tables 3-5 and Figure 1a-e. ISSR bands were treated as there or absent, exclusive of considering their percentage. The middling number of augmented loci and the middling number of polymorphic loci per primer were 11.6 and 2.6, respectively, as the percent of loci polymorphism ranged from 0.0% to 58.0% with an average of 21.4% crossways all the flax genotypes. An entirety of 139 dissimilar fragments have been recorded and formed mainly by six of the 12 used primers, showing 14 bands by primer ISSR-2 ranging from 1.15 to 0.15 Kbp, 12 bands by primer ISSR-5 ranging from 0.85 to 0.17 Kbp, and 11 bands by primer ISSR-3 and ISSR-10 ranging from 1.20 to 0.21 and 1.45, 0.70 Kbp, respectively. On the other hand, primers ISSR-1, ISSR-4, and ISSR-8 produced 10 bands. Moreover, primers R-5, R-6, and ISSR-6 produced only 7 and 6 bands, respectively. The analysis of ISSR-PCR data can thus select the use of primers ISSR-2, ISSR-5, ISSR-3, and ISSR-10 for the selective discrimination of flax genotypes from other commercial varieties. These primers may be used as an indicator for obtaining genetic markers. Out of 139 loci detected, the polymorphic, monomorphic, and unique loci were 31, 108, and 13, respectively. All flax cultivars were discriminated by the presence or absence of unique piece in ISSR outline.

Table 2: Mean values of straw, seed, fiber, and oil yields, as well as fiber and oil percentages for five flax genotypes from the combined analysis over the two seasons (2014/2015) and (2015/2016)

Genotype	Straw (ton/feddan)	Seed (kg/feddan)	Fiber (kg/feddan)	Oil (kg/feddan)	Fiber (%)	Oil (%)
Sakha 1	3.65	630.00	597.14	250.11	16.36	39.70
S.533/39/5/3 (F1)	4.70	470.00	814.35	174.37	18.30	37.10
S.402/3/3/7 (F2)	3.78	725.00	650.09	294.35	17.22	40.60
S.421/3/6/4 (F3)	3.95	743.00	687.30	304.85	17.40	41.03
S.11 (F4)	4.06	430.00	730.80	157.81	18.00	36.70

These results are means of four replications

Cluster analysis

A dendrogram was generated based on the resemblance matrix by the UPGMA, in which the flax genotypes were grouped in three clusters. The Jaccard's similarity coefficient among the genotypes ranged from 0.91 to 0.95 [Figure 2].

DISCUSSION

To go for a proper conservation, management, and selection of parental lines for large-scale cultivation, a prior knowledge on the existing genetic diversity is most important. For improvement of crop genetic resource, it is necessary to have continuous mixing of wild relatives and use of effective breeding methods. Considering the need of molecular characterization of the crop, studies were undertaken on DNA fingerprinting using ISSR and RAPD markers.^[27]

Flax is a highly important crop valued for its fixed oil and fibers which are used in different industries, in addition to the medicinal importance of its lignans. To enrich the genetic pool of the species, screening and selection of genotypes are essential, and this can be only achieved through proper genetic diversity study. Several earlier works were done on different flax cultivars, but this is the first attempt to fingerprint the four promising flax genotypes derived from the main commercial variety Sakha 1 which is cultivated in Egypt. The DNAs from the fresh leaves of the four flax genotypes have been compared. Study of the degree and

allotment of genetic diversity in harvest vegetation was performed for optimizing variety and breeding strategies.

ISSRs are arbitrary multilocus markers produced by PCR amplification with a single anchored microsatellite primer. They are beneficial because no genomic in order is necessary for their employ. This provides a suitable and rapid appraisal of the differences in the genetic composition of closely related individuals at the DNA level and has been working in a large number of plant species for categorization and estimation of genetic assortment because of their pace and simplicity in handling.^[28] ISSR markers were successfully applied to evaluate genetic diversity and relations among 22 Canadian cultivars, 29 selected world cultivars and 10 landraces of flax.^[9,29]

In the present research, ISSR analysis made it possible to detect the changeability in the majority of loci of different parts of the genome and obtain the objective evidence about the genetic relations between flax genotypes. The four promising genotypes of flax were subjected to realized analysis with the help of 12 effective random primers, ISSR-1, ISSR-2, ISSR-3, ISSR-4, ISSR-5, ISSR-6, ISSR-7, ISSR-8, ISSR-9, ISSR-10, R-5, and R-6. The number of ISSR-PCR fragments indicated that the 12 primers were reproduced. Each DNA band was treated as a unit character. There were 139 loci obtained randomly and distributed over genome and 108 common for all cultivars fragments of amplification, as well as 31 polymorphic fragments. The three primers 5'-GAC (GATA)₄, 5'-(GATA)₄GC, and 5'-(AGAC)₄GC exhibited 58%, 50%, and 44% polymorphism.

The high level of polymorphism observed in this study indicated a elevated level of genetic variation among the 4 genotypes analyzed, these results were in accordance with Rakoczy-Trojanowska and Bolibok,^[30] who reported highly polymorphic patterns when retort primers based on microsatellite sequences in plants were working. Our work was also in accordance with that reported by Blair *et al.*,^[31] where primers with poly-GA motifs produce on average, a larger number of amplified loci. In the nearby effort, primers with 5' anchoring showed more monomorphic loci (seven from 12 primers) [Table S1]. These outcomes could be doable because primers anchoring in 5' include in the amplified creation the whole microsatellite sequence, and thus, the variability in the number of nucleotides inside a microsatellite repeat would consequence in length polymorphism when using a 5' anchored primer.^[32]

The dendrogram obtained by the UPGMA method allowed the identification of two major clusters. The first one comprised F1 and F2 which had 95% similarity, whereas the subsequent comprised F₁, F₂, and F₄ which had 94% resemblance to each other. On the extra hand

Table 3: The sequence and names of inter simple sequence repeat primers used in the fingerprinting of the five Egyptian cultivars

Name	Sequence	Primer	T° (°C)
ISSR-1	5'-AGAGAGAGAGAGAGAGC-3'	(AG) ₈ C	50
ISSR-2	5'-AGAGAGAGAGAGAGAGG-3'	(AG) ₈ G	50
ISSR-3	5'-ACACACACACACACACT-3'	(AC) ₈ T	51
ISSR-4	5'-ACACACACACACACACG-3'	(AC) ₈ G	53
ISSR-5	5'-GTGTGTGTGTGTGTG-3'	(GT) ₈ G	51
ISSR-6	5'-CGCGATAGATAGATAGATA-3'	CGC (GATA) ₄	49
ISSR-7	5'-GACGATAGATAGATAGATA-3'	GAC (GATA) ₄	58
ISSR-8	5'-AGACAGACAGACAGACGC-3'	(AGAC) ₄ GC	55
ISSR-9	5'-GATAGATAGATAGATAGC-3'	(GATA) ₄ GC	50
ISSR-10	5'-GACAGACAGACAGACAAT-3'	(GACA) ₄ AT	49
R-5	5'-ACACACACACACACACA-3'	(AC) ₈ A	
R-6	5'-ACACACACACACACACC-3'	(AC) ₈ C	

A: Adenine; T: Thymine; G: Guanine; C: Cytosine; T°: Annealing temperature; °C: Centigrade degree; ISSR: Inter simple sequence repeat

Table 4: Degree of polymorphism and polymorphic information content for interspecies genetic relationship in 4 flax cultivars*

Primers	Gel polymorphism						
	TL	ML	PL polymorphic (without unique)	Unique bands	Polymorphic (with unique)	Polymorphism (%)	Mean of band frequency
(AG) ₈ C	12	10	0	2	2	17	0.9
(AG) ₈ G	15	14	1	0	1	7	1.0
(AC) ₈ T	11	11	0	0	0	0	1.0
(AC) ₈ G	11	10	0	1	1	9	0.9
(GT) ₈ G	12	12	0	0	0	0	1.0
CGC (GATA) ₄	8	6	0	2	2	25	0.8
GAC (GATA) ₄	12	5	6	1	7	58	0.7
(AGAC) ₄ GC	18	10	5	3	8	44	0.8
(GATA) ₄ GC	10	5	3	2	5	50	0.8
(GACA) ₄ AT	14	11	2	1	3	21	0.9
(AC) ₈ A	8	7	0	1	1	13	0.9
(AC) ₈ C	8	7	1	0	1	13	0.9
Total	139	108	18	13	31		
Mean	11.58	9	1.5	1.08	2.58	21.41	

*Flax genotypes: Namely, promising strains (F1) 533/39/5/3, (F2) S.402/3/3/7, (F3) S.421/3/6/4, and (F4) S.11. TL: Total loci; ML: Monomorphic loci; PL: Polymorphic loci; PIC: Polymorphic information content




Figure 1: (a) Inter simple sequence repeat – polymerase chain reaction profiles were obtained using a primers inter simple sequence repeat 1, inter simple sequence repeat 2, and inter simple sequence repeat 3 in different flax cultivars F1=, F2=, F3=, F4 = respectively; (lanes: 1-4)=, Lane M = DNA ladder. Arrows show cultivars specific band. (b) inter simple sequence repeat – polymerase chain reaction profiles were obtained using a primers inter simple sequence repeat 4, inter simple sequence repeat 4, and inter simple sequence repeat 5 F1=, F2=, F3=, F4 = respectively; (lanes: 1-4)=, Lane M = DNA ladder. (c) inter simple sequence repeat – polymerase chain reaction profiles were obtained using a primers inter simple sequence repeat 7 and inter simple sequence repeat 8 F1=, F2=, F3=, F4 = respectively; (lanes: 1-4)=, Lane M = DNA ladder. (d) inter simple sequence repeat – polymerase chain reaction profiles were obtained using a primers inter simple sequence repeat 9 and inter simple sequence repeat 10 F1=, F2=, F3=, F4 = respectively; (lanes: 1-4)=, Lane M = DNA ladder. (e) Inter simple sequence repeat analysis carried out with primers R5 and R6 F1=, F2=, F3=, F4 = respectively; (lanes: 1-4)=, Lane M = DNA ladder

over, the third group comprised F3, which had 91% similarity to the second group.

The phenotypic variations of the four flax genotypes on the basis of the straw, seed, fiber, and oil yield have also supported the segregation among them. Where, F3 genotype exhibited the highest seed yield (743 Kg/fed.) as well as oil yield (304.85 Kg/fed.) and oil percent (41.03%). Industrially, flaxseed oil is an important ingredient in the manufacture of paint, varnish, and linoleum.^[17] Flaxseed oil also contains α -linolenic acid, a polyunsaturated fatty acid that has nutritional and health benefits


(Wood, 1997).^[18] Our study therefore identified F3 to be the major assorted genotypes and recommended their use in breeding programs and for upward mapping populations.

The incidence of environmentally induced transmissible changes in definite flax varieties has been shown to be accompanied by changes in the genomic DNA.^[33] Introduction of modern cultivars and using them in breeding is the only way to ensure the decrease in genetic variation.^[34,35] Shrinking of the genetic diversity consequently reduces options to ensure diverse nutrition, to enhance food production and to face climate

Table 5: Genetic similarity matrix based on inter simple sequence repeat – polymerase chain reaction data among* four flax genotypes (F1–F4) estimated according to Jaccard's method

	Similarity matrix			
	F1	F2	F3	F4
F1	100			
F2	95	100		
F3	91	91	100	
F4	94	95	91	100

*Flax genotypes: Namely, promising strains (F1) 533/39/5/3, (F2) S.402/3/3/7, (F3) S.421/3/6/4 and (F4) S.11

**Figure 2:** Unweighted pair-group method with arithmetic average dendrogram showing clustering of 4 flax cultivars based on inter simple sequence repeat – polymerase chain reaction data

change. Crop wild relatives hosted in gene bank collections have received some attention because they harbor untapped genetic variation available for domesticated crops.^[36,37] Despite the few marker-trait associations identified, herein, we provided a proof of thought for brown flax functional variants as potentially useful for cultivated flax improvement.

In the present study, we demonstrated that the ISSR primers can be effectively working to evaluate the level of polymorphism and diversity in flax genotypes. The results obtained in this study legalize once more that ISSRs are useful markers in genetic variety studies, due to the very high polymorphism level detected by the primers. The possibility of classification of every individual examined offers a promising perception as a molecular tool for varietal recognition and breeding program applications.

CONCLUSION

To the best of our awareness, this is the first exertion to judge genetic relationships among the Egyptian flax genotypes and to use molecular markers for categorization of the influential flax genotypes actively concerned in Egyptian flax breeding programs. These findings will be certainly helpful for the breeders in the selection of genotypes for the future breeding and improvement programs.

Supporting information

Data of the unique and common bands for the four flax cultivars obtained with 12 primers in PCR amplification is available as supporting information.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Zohary D. Monophyletic and polphylectic origin of the crops on which agriculture was formed in the Near East. *Genet Resour Crop Evol* 1999;46:133-42.
- Millam S, Obert B, Pretova A. Plant cell and biotechnology studies in *Linum usitatissimum* (a review). *Plant Cell Tissue Organ Cult* 2005;82:93-103.
- Carter JF. Potential of flaxseed and flax oil in baked goods and other products in human nutrition. *Cereal Food World* 1993;38:753-9.
- Oh TJ, Gorman M, Cullis CA. RAPD mapping in flax (*Linum usitatissimum* L.). *Theor Appl* 2000;101:590-3.
- Mansby E, Diaz O, von Bothmer R. Preliminary study of genetic diversity in Swedish flax (*Linum usitatissimum*). *Genet Resour Crop Evol* 2000;47:417-24.
- Diederichsen A, Richards K. Cultivated flax and the genus *Linum* L. taxonomy and germplasm conservation. In: Muir AD, Westcott ND, editors. *Flax: The Genus Linum*. Boca Raton: CRC Press; 2003. p. 22-54.
- Miah G, Rafii MY, Ismail MR, Puteh AB, Rahim HA, Islam KhN, *et al.* A review of microsatellite markers and their applications in rice breeding programs to improve blast disease resistance. *Int J Mol Sci* 2013;14:22499-528.
- Yashitola J, Thirumurugan RM, Sundaram MK, Naseerullah MS, Ramesha MS, Sarma NP. Assessment of purity of rice hybrids using microsatellite and STS markers. *Crop Sci* 2002;42:1369-73.
- Fu Y, Diederichsen A, Richards K. Genetic diversity within a range of cultivars and landraces of flax (*Linum usitatissimum* L.) as revealed by RAPDs. *Genet Resour Crop Evol* 2002;49:167-7.
- Fu YB, Rowland GG, Duguid SD, Richards KW. RAPD analysis of 54 North American flax cultivars. *Crop Sci* 2003;43:1510-5.
- Diederichsen A, Fu YB. Phenotypic and molecular (RAPD) differentiation of four infraspecific groups of cultivated flax (*Linum usitatissimum* L. subsp *usitatissimum*). *Genet Resour Crop Evol* 2006;53:77-90.
- Wiesner I, Wiesnerová D. Insertion of a reamplification round into the ISSR-PCR protocol gives new flax fingerprinting patterns. *Cell Mol Biol Lett* 2003;8:743-8.
- Wiesnerová D, Wiesner I. ISSR-based clustering of cultivated flax germplasm is statistically correlated to thousand seed mass. *Mol Biotechnol* 2004;26:207-14.
- Rajwade AV, Arora RS, Kadoo NY, Harsulkar AM, Ghorpade PB, Gupta VS, *et al.* Relatedness of Indian flax genotypes (*Linum usitatissimum* L.): An inter-simple sequence repeat (ISSR) primer assay. *Mol Biotechnol* 2010;45:161-70.
- Uysal H, Fu YB, Kurt O, Peterson GW, Diederichsen A, Kusters P. Genetic diversity of cultivated flax (*Linum usitatissimum* L.) and its wild progenitor pale flax (*Linum bienne* Mill.) as revealed by ISSR markers. *Genet Resour Crop Evol* 2010;57:1109-19.
- Morris D.H. Flax: A health and nutrition primer. Winnipeg: Flax Council of Canada; 2003. p. 11.
- Matheson EM. Linseed. In: *Vegetable Oil Seed Crops in Australia*. Sydney: Holt, Rinehart and Winston; 1976. p. 111-21.
- Wood IM. Fibre Crops – New Opportunities for Australian Agriculture. Brisbane: Department of Primary Industries; 1997. p. 18-24.
- Easson DL, Molloy RM. A study of the plant, fibre and seed development in flax and linseed (*Linum usitatissimum* L.) grown at a range of seed rates. *J Agric Sci* 2000;135:361-9.
- Abd El-Mohsen AA, Abdallah AM, Mahmoud GO. Optimizing and describing the influence of planting dates and seeding rates on flax cultivars under Middle Egypt region conditions. *World Essays J* 2013;12:28-39.
- Doyle JJ, Doyle JL. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull* 1987;19:11-1.
- Poyraz İ. An efficient DNA isolation method from *Nigella sativa* L. (Ranunculaceae) seeds for RAPD and ISSR analysis. *Bilecik Sheikh Edebalı University Science Journal* 2014;1:22-7.
- Williams JG, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res* 1990;18:6531-5.
- Sneath PH, Sokal RR. *Numerical Taxonomy*. San Francisco, CA, USA: Freeman Press; 1973.
- Snedecor GW, Cochran WG. *Statistical Method*. 7th ed. Ames: Iowa State University Press; 1982. p. 325-30.
- Le Clerg E, Leonard WE, Clark AG. *Field Plot Technique*. Minneapolis,

- Minnesota: Burgross Publishing Co.; 1966.
27. Baruah J, Gogoi B, Das K, Ahmed NM, Sarmah DK, La M, *et al.* Genetic diversity study amongst *Cymbopogon* species from NE-India using RAPD and ISSR markers. *Ind Crops Prod* 2017;95:23-43.
 28. Zhao WG, Miao XX, Zang B, Zhang L, Pan YL, Huang YP, *et al.* Construction of fingerprinting and genetic diversity of mulberry cultivars in China by ISSR markers. *Yi Chuan Xue Bao* 2006;33:851-60.
 29. Kenaschuk EO, Rowland GG. Flax. In: Slinkard AE, Knott DR, editors. *Harvest of Gold: The History of Field Crop Breeding in Canada*. Canada: University of Saskatchewan SK; 1995. p. 173-6.
 30. Rakoczy-Trojanowska M, Bolibok H. Characteristics and a comparison of three classes of microsatellite-based markers and their application in plants. *Cell Mol Biol Lett* 2004;9:221-38.
 31. Blair MW, Panaud O, McCouch SR. Inter-simple sequence repeat (ISSR) amplification for analysis of microsatellite motif frequency and fingerprinting in rice (*Oryza sativa* L.). *Theor Appl Genet* 1999;98:780-92.
 32. Pradeep Reddy M, Sarla N, Siddiq EA. Inter simple sequence repeat (ISSR) polymorphism and its application in plant breeding. *Euphytica* 2002;128:9-17.
 33. Cullis CA, Swami S, Song Y. RAPD polymorphisms detected among the flax genotypes. *Plant Mol Biol* 1999;41:795-800.
 34. Tanksley SD, McCouch SR. Seed banks and molecular maps: Unlocking genetic potential from the wild. *Science* 1997;277:1063-6.
 35. Vromans J. *Molecular Genetic Studies in Flax (Linum usitatissimum L.)*. Ph. D. Thesis, Wageningen University, The Netherlands; 2006.
 36. McCouch S, Baute GJ, Bradeen J, Bramel P, Bretting PK, Buckler E, *et al.* Agriculture: Feeding the future. *Nature* 2013;499:23-4.
 37. Xiao J, Li J, Grandillo S, Ahn SN, Yuan L, Tanksley SD, *et al.* Identification of trait-improving quantitative trait loci alleles from a wild rice relative, *Oryza rufipogon*. *Genetics* 1998;150:899-909.

Table S1: The unique and common bands for flax genotypes (F1–F4) obtained with 12 decamer primers in polymerase chain reaction amplification

Primers	Cultivars (specific unique bands [bp])			
	F1	F2	F3	F4
ISSR-1 [Figure 2a]	1500	1500	1500	1500
	1300	1300	1300	1300
	1050	1050	1050	1050
	810	810	810	810
	780	780	780	780
	660	660	660	660
	-	590	-	-
	540	540	540	540
	-	510	-	-
	440	440	440	440
ISSR-2 [Figure 2a]	340	340	340	340
	220	220	220	220
	1150	1150	1150	1150
	1000	1000	1000	1000
	790	790	790	790
	690	-	-	690
	590	590	590	590
	510	510	510	510
	480	480	480	480
	420	420	420	420
ISSR-3 [Figure 2a]	360	360	360	360
	320	320	320	320
	300	300	300	300
	280	280	280	280
	230	230	230	230
	210	210	210	210
	150	150	150	150
	1200	1200	1200	1200
	940	940	940	940
	880	880	880	880
ISSR-4 [Figure 2b]	730	730	730	730
	620	620	620	620
	580	580	580	580
	520	520	520	520
	420	420	420	420
	290	290	290	290
	270	270	270	270
	210	210	210	210
	950	950	950	950
	790	790	790	790
ISSR-5 [Figure 2b]	700	700	700	700
	620	620	620	620
	560	560	560	560
	500	500	500	500
	440	440	440	440
	410	410	410	410
	-	-	370	-
	280	280	280	280
	260	260	260	260
	850	850	850	850
ISSR-6 [Figure 2b]	730	730	730	730
	680	680	680	680
	510	510	510	510
	480	480	480	480
	430	430	430	430
	370	370	370	370
	330	330	330	330
	300	300	300	300
	270	270	270	270
	170	170	170	170

Contd...

Table S1: Contd...

Primers	Cultivars (specific unique bands [bp])			
	F1	F2	F3	F4
ISSR-6 [Figure 2b]	-	1250	-	-
	790	790	790	790
	710	710	710	710
	600	600	600	600
	490	490	490	490
	460	460	460	460
	410	-	-	-
	390	390	390	390
	-	1250	1250	-
	-	-	1000	-
ISSR-7 [Figure 2c]	850	-	850	-
	790	790	790	790
	710	710	710	710
	580	580	580	580
	480	480	480	480
	440	440	440	440
	400	400	-	-
	-	380	380	380
	320	320	-	-
	-	290	290	290
ISSR-8 [Figure 2d]	1150	1150	-	1150
	-	-	960	-
	900	900	-	900
	860	860	860	860
	660	660	660	660
	640	640	640	640
	560	560	560	560
	460	460	-	460
	450	450	450	450
	-	-	440	-
ISSR-9 [Figure 2d]	410	410	410	410
	380	380	380	380
	350	350	350	350
	310	310	310	310
	-	-	280	-
	260	260	260	260
	230	230	-	-
	200	-	200	-
	-	-	1400	-
	1200	1200	1200	1200
ISSR-10 [Figure 2d]	1000	1000	1000	1000
	810	810	810	810
	700	700	700	700
	620	620	620	-
	-	-	-	570
	550	550	550	550
	510	-	510	-
	380	380	380	-
	1450	1450	1450	1450
	-	-	1150	-
ISSR-11 [Figure 2d]	1050	1050	1050	1050
	950	950	950	950
	810	810	-	810
	-	750	-	750
	700	700	700	700
	590	590	590	590
	550	550	550	550
	470	470	470	470
	420	420	420	420
	370	370	370	370
ISSR-12 [Figure 2d]	290	290	290	290
	170	170	170	170

Contd...

Table S1: Contd...

Primers	Cultivars (specific unique bands [bp])			
	F1	F2	F3	F4
R5 [Figure 2e]	1350	1350	1350	1350
	1050	1050	1050	1050
	870	870	870	870
	680	565	565	565
	580	580	580	580
	-	-	500	-
	380	433	433	433
	250	369	369	369
	820	820	820	820
	600	600	600	600
R6 [Figure 2e]	550	550	550	550
	380	380	380	380
	340	340	340	340
	-	-	320	320
	300	300	300	300
	250	250	250	250

Flax genotypes: Namely, promising strains (F1) 533/39/5/3, (F2) S.402/3/3/7, (F3) S.421/3/6/4 and (F4) S.11. ISSR: Inter simple sequence repeat