

Triterpenes from *Euphorbia rigida*

Noureddine Gherraf, Amar Zellaoui, Naglaa S. Mohamed¹, Taha A. Hussien², Tarik A. Mohamed³, Mohamed-Elamir F. Hegazy⁴, Salah Rhouati, Mahmoud F. M. Moustafa⁵, Magdi A. El-Sayed⁶, Abou El-Hamd H. Mohamed¹

Laboratory of Natural Products and Organic Synthesis, Department of Chemistry, Faculty of Science, University Mentouri - Constantine, Algeria, Departments of ¹Chemistry, ⁶Botany, Aswan-Faculty of Science, South Valley University, Aswan, Egypt, ²Department of Pharmaceutical Chemistry and Pharmacognosy, Faculty of Pharmacy, Applied Sciences University, Amman, Jordan - 119 31, ³Department of Pharmacognosy, Faculty of Pharmacy, El-Minia University, El-Minia - 615 19, ⁴Chemistry of Medicinal Plant Department, National Research Centre, Dokki, Giza - 126 22, Egypt, ⁵Department of Biological Sciences, College of Science, King Khalid University, 61413, Saudi Arabia

Submitted: 20-02-2010

Revised: 05-06-2010

Published: 19-07-2010

ABSTRACT

Phytochemical studies of the aerial parts of *Euphorbia rigida* afforded three triterpenes: betulin (1), cycloart-23Z-ene-3, 25-diol (2) and cycloartan-3, 24, 25-triol (3), firstly isolated from this plant. The structures and relative stereochemistry were determined on the basis of extensive spectroscopic analyses, including 1D and 2D NMR experiments (1H NMR, 13C NMR, COSY, NOESY, HMQC and HMBC).

Key words: *Euphorbia rigida*, Euphorbiaceae, cycloartan triterpene

INTRODUCTION

Euphorbia genus belongs to the family Euphorbiaceae. This family comprises about 300 genus and 5000 species distributed mainly in America and tropical Africa.^[1] *Euphorbia* species have been used in folk medicine to treat skin diseases, migraines, intestinal parasites and warts.^[2] The biological activities of the genus, including antitumor, antiviral, cytotoxic properties and different vascular effects, are generally attributed to the presence of specific types of diterpenes, both macrocyclic and polycyclic derivatives.^[3-5] The skin irritant and tumor-promoting properties of tiglane, ingenane and dephanane diterpenes of this plant are well known. Considerable attention has recently been given to the macrocyclic diterpenes because of their high chemical diversity and therapeutically relevant bioactivity.^[6-8] Jatrophone and modified jatrophone diterpenoids, which are rare in the genus *Euphorbia*, are potent inhibitors of a membrane protein (so-called P-glycoprotein) pumping cytotoxic drugs out of cells and conferring upon the cells the ability to resist high doses of these drugs.^[9] Therefore, the genus has been subjected to numerous chemical studies and these have led to the isolation of diterpenes^[10,11] dimeric

diterpenoid^[12] diterpene polyesters^[11,13] triterpenes^[14] and pentacyclic triterpenes.^[15] Few sesquiterpenoids and flavonoids have been isolated from the genus.^[16,17]

Spurges *Euphorbia* species are a common constituent of many ancient treatments of mouse leukemia and diseases such as cancer, swelling and warts.^[18]

MATERIALS AND METHODS

Plant material

The aerial parts of *Euphorbia rigida* were collected from Greek in August 2004, by Dr. Olga Tzakou, Department of pharmacy and Chemistry of Natural products, Faculty of Pharmacy, University of Athens, Greece.

Extraction and isolation

The air-dried plant (1 kg) was crushed and extracted with CH₂Cl₂-MeOH (1:1) at room temperature. The extract was concentrated *in vacuo* to obtain a residue (30 g). The residue was fractionated by silica gel CC (6 × 120 cm) eluted with *n*-hexane (3 l), followed by a gradient of *n*-hexane-CH₂Cl₂ up to 100% CH₂Cl₂ and CH₂Cl₂-MeOH up to 15% MeOH (2 l of each solvent mixture) with increasing degree of polarity. The *n*-hexane-CH₂Cl₂ (1:1) was pre-fractionated by CC using Sephadex LH-20 (2 × 40 cm) and eluted with *n*-hexane-CH₂Cl₂ (7:4) to give compound **1** (80 mg). Compound **2** (60 mg) was isolated from *n*-hexane-CH₂Cl₂ (2:3) fraction and the latex was pre-fractionated by CC on Sephadex LH-20 (1 × 30 cm) and eluted with *n*-hexane-

Address for correspondence:

Dr. Magdi A. El-Sayed,
Department of Botany, Aswan-Faculty of Science,
South Valley University, Aswan, Egypt.
E-mail: magradi2000@yahoo.com

DOI: 10.4103/0974-8490.65510

CH₂Cl₂-MeOH (7:4:0.25). Compound **3** (30 mg) isolated from 100% CH₂Cl₂ fraction, was pre-fractionated by CC on Sephadex LH-20 (1 × 30 cm) and eluted with *n*-hexane-CH₂Cl₂-MeOH (7:4:0.5).

RESULTS AND DISCUSSION

Repetitive chromatographic steps of the methylenechloride/methanol (1:1) extract of the air-dried aerial parts of *E. rigida* yielded three known triterpenes [Figure 1].

Compound **1** was obtained as a white powder. The structure of **1** was determined from careful investigation of the 1D and 2D NMR measurements. The ¹H-NMR spectrum (600 MHz, CDCl₃) [Table 1] showed the triterpenoid pattern with six methyl singlets in the up-field at δ_H 0.74 (Me-24), 0.79 (Me-25), 0.92 (Me-23), 0.94 (Me-27), 0.99 (Me-26) and the one methyl group of Me-30 appeared as a sharp singlet at δ_H 1.68 (Me-30). The down-field shift for Me-30 indicated the presence of a double bond between C-20 and C-29. In the down-field of spectrum, there were two broad singlets: at δ_H 4.65 (1H, br s, H-29_a) and 4.55 (1H, br s, H-29_b), suggesting the presence of an olefinic proton. The HMQC spectrum showed correlations between H-29_a at δ_H 4.65 (1H, br s) and H-29_b at δ_H 4.55 (1H, br s) with carbon signal at δ_C 109.72. Additionally, the hydroxylated methylene protons at δ_H 3.75 (1H, d, *J* = 9 Hz, H-28_a), coupled in ¹H-¹H COSY spectrum with a signal at δ_H 3.32 (1H, d, *J* = 9 Hz, H-28_b), giving a doublet. The HMQC spectrum showed correlation between H28_a at δ_H 3.75 (1H, d, *J* = 9 Hz) and H-28_b at δ_H 3.32 (1H, d, *J* = 9 Hz) with carbon at δ_C 60.39. The ¹H NMR also revealed a secondary

hydroxyl group placed at C-3, inferred from the down-field shift of methine proton which appeared at δ_H 3.18 (1H, dd, *J* = 8, 3.2 Hz, H-3) which showed correlation in HMQC with carbon signal at δ_C 78.86.

The ¹³C-NMR spectrum (125 MHz, CDCl₃) [Table 2] displayed 30 carbon signals and DEPT experiments indicated these signals corresponding to 6 methyl groups, 12 methylene groups, including one attached to oxygen appearing at δ_C 60.42 for C-28. Six methine groups including one attached to oxygen appeared at δ_C 78.86 for C-3 and six quaternary carbon atoms. The olefinic carbons C-20 and C-29 appeared at δ_C 15.46 and 109.77, respectively. HMQC and HMBC were used to determine the position of the hydroxylated methyl carbons; the two proton signals at δ_H 3.75 (H-28_a) and 3.32 (H-28_b) seen in the HMBC experiment show clear

Table 1: ¹H NMR spectroscopic data for compounds 1–3 (600 MHz, CDCl₃)

Position	1	2	3
H-1	1.65 dd (3.6, 12.6) 0.89 dd (3.6, 12.6)	1.30 m 0.90 m	1.60 m
H-2	1.58 m	1.4 m 1.23 m	1.65 m
H-3	3.18 dd (3.2, 8)	3.10 m	3.35 t (3.2)
H-5	0.33 br d (18)	0.91 m	1.25 m
H-6	1.50 m 1.38 m	1.26 m 0.45 dd (12.5, 8)	1.50 m
H-7	1.37 m	1.65 m 1.40 m	1.72 m
H-8		1.16 dd (5,13)	1.16 dd (3, 8)
H-9	1.29 m		
H-11	1.40 m 1.19 m	0.77 m	0.11 m
H-12	1.62 m 1.05 m	0.93 m	1.30 m
H-13	1.63 m		
H-15	1.63 m 1.69 m	1.27 m	0.15 m
H-16	1.89 m 1.28 m	1.60 m	1.88 m
H-17		1.24 m	1.50 m
H-18	1.58 m	0.62 s	0.97 s
H-19	1.36 m	0.22 d (4.5) 0.01d (4.5)	0.53 d (4.5) 0.30 d (4.5)
H-20		1.10 m	1.35 m
H-21	1.98 m	0.53 d (3.5)	0.87 d (3.2)
H-22	1.29 m 1.03 m 1.85 m	1.88 m	2.22 m
H-23	0.92 s	5.26 d (8)	3.10 m
H-24	0.74 s	5.26 d (8)	3.25 m
H-25	0.79 s		
H-26	0.99 s	0.98 s	1.12 s
H-27	0.94 s	0.98 s	1.24 s
H-28	3.30 d (9) 3.75 d (9)	0.62 s	0.97 s
H-29	4.65 br s 4.55 br s	0.42 s	0.75 s
H-30	1.68 s	0.56 s	0.88 s




Figure 1: Selected HMBC correlations of compound **1**

Table 2: ^{13}C NMR spectroscopic data for compounds 1–3 (125 MHz, CDCl_3)^a

Position	1	2	3
C-1	38.64 t	31.93 t	31.97 t
C-2	27.03 t	30.33 t	30.38 t
C-3	78.86 d	78.80 d	78.83 d
C-4	38.82 s	40.45 s	40.48 s
C-5	55.26 d	47.10 d	47.11 d
C-6	18.28 t	21.08 t	21.11 t
C-7	34.21 t	28.04 t	28.36 t
C-8	40.89 s	48.00 d	47.97 d
C-9	50.36 d	20.00 s	19.93 s
C-10	37.13 s	25.96 s	26.00 s
C-11	20.80 t	26.07 t	26.02 t
C-12	25.17 t	35.55 t	35.54 t
C-13	37.28 d	45.29 s	45.42 s
C-14	42.69 s	48.80 s	48.83 s
C-15	27.01 t	32.76 t	33.01 t
C-16	29.15 t	26.42 t	26.46 t
C-17	47.74 s	52.00 d	52.29 d
C-18	48.72 d	18.04 q	17.99 q
C-19	47.77 d	30.00 t	29.88 t
C-20	150.46 s	36.36 d	36.35 d
C-21	29.72 t	18.25 q	18.12 q
C-22	33.95 t	39.00 t	33.31 t
C-23	27.96 q	139.31 d	28.41 t
C-24	15.35 q	125.57 d	79.63 d
C-25	16.08 q	70.75 s	76.74 s
C-26	15.95 q	29.66 q	23.24 q
C-27	14.74 q	29.83 q	26.54 q
C-28	60.42 t	19.26 q	19.26 q
C-29	109.72 t	13.9 q	13.98 q
C-30	19.06 q	25.41 q	25.42 q

^aMultiplicity was determined by DEPT experiments (s, quaternary; d, methine; t, methylene; q, methyl)

long-range correlations between the carbon signals at δ_{C} 29.15 (C-16), 33.95 (C-22) and 47.74 (C-17), while the carbon signal at δ_{C} 60.39 (C-28) showed a correlation with the proton signal at δ_{H} 1.03 (H-22_a), 1.85 (H-22_b), 1.28 (H-16_a). Other important correlations were observed between the carbon signals at δ_{C} 15.35 (C-24), 27.96 (C-23) and 38.64 (C-1) with the proton signal at δ_{H} 3.18 (H-3). Therefore, the hydroxylated methyl was placed at C-3. The assignment of all proton signals and their connectivity to adjacent protons and carbon signals were established from the results of 2D ^1H - ^1H COSY and HMQC experiments.

Acetylation of **1** gave the diacetyl derivative (**1a**), for which the ^1H NMR spectrum displayed two new acetyl signals and confirmed the structure of compound **1**. The structure of compound **1** was deduced from the comparison of its spectral data with those of literature and identified as betulin.^[19,20]

Compound **2** was isolated as colorless needles. The ^1H -NMR spectrum (600 MHz, CDCl_3) [Table 1] of compound **2** displayed two doublets at δ_{H} 0.22 (1H, d, $J = 4.5$ Hz,

H-19a) and at 0.01 (1H, d, $J = 4.5$ Hz, H-19b) which are characteristic of the presence of a C-9/C-10 cyclopropyl methylene group of a cycloartan-3-one triterpenoid. Cycloartane-type triterpenes possess a cyclopropane bridge between C-9 and C-10, and protons attached to cyclopropyl rings characteristically appear as a pair of doublets in the high-field ^1H -NMR region with gem-coupling constant ($J = 4.5$ Hz). The ^1H -NMR spectrum showed the presence of an olefinic proton double bond at δ_{H} 5.26 (2H, d, $J = 8$, H-23, H-24). The low coupling constant ($J = 8$ Hz) between H-23 and H-24 indicate the stereochemistry “Z” at C-23. The HMQC spectrum showed correlation between H-23 at δ_{H} 5.26 (1H, d, $J = 8$) with carbon signal at δ_{C} 139.31 and H-24 at δ_{H} 5.26 (1H, d, $J = 8$) with carbon signal at δ_{C} 125.57. Additionally, δ_{H} 2.95 (1H, m, H-3) which suggested the existence of secondary hydroxyl group. The ^1H -NMR spectrum showed the presence of six tertiary methyl groups as a singlet at δ_{H} 0.42 (3H, s, Me-29), 0.56 (3H, s, Me-30), a sharp signal that integrated for six protons at δ_{H} 0.62 (6H, s, Me-18, Me-28), 0.98 (6H, s, Me-26, Me-27) and one methyl group at δ_{H} 0.53 (3H, d, $J = 6.5$ Hz, Me-21) coupled with H-20 (methine proton) gave a doublet.

The ^{13}C -NMR spectrum (125 MHz, CDCl_3) [Table 2] of compound **2** showed the presence of 30 carbon signals. Determination of the multiplicity was carried out by DEPT experiments which indicated the presence of 6 quaternary carbon atoms, 7 methine groups, 10 methylene groups and 7 methyl groups. It also showed the presence of two olefinic carbons C-23 and C-24 appearing at δ_{C} 139.31 and 125.57, respectively. Two oxygenated carbons appeared at δ_{C} 78.80 for C-3 and at 70.75 for C-25. The structure of compound **2** was deduced from the comparison of its spectral data with those of literature and identified as cycloart-23Z-ene-3, 25-diol.^[21,22]

Compound **3** was isolated as colorless needles. The ^1H -NMR spectrum (500 MHz, CDCl_3) [Table 1] of compound **3** displayed two doublets at δ_{H} 0.55 (1H, d, $J = 4.5$ Hz, H-19_a) and at 0.3 (1H, d, $J = 4.5$ Hz, H-19_b) which are characteristic of a C-9/C-10 cyclopropyl methylene group of a cycloartan-3-one triterpenoid. Cycloartane-type triterpenes possess a cyclopropane bridge between C-9 and C-10, and protons attached to cyclopropyl rings characteristically appear as a pair of doublets in the high-field ^1H -NMR region with a gem-coupling constant ($J = 4.5$ Hz). Additionally, the presence of triplet bonds at δ_{H} 3.35 (1H, t, $J = 3.2$ Hz, H-3) and multiplet bands at δ_{H} 3.25 (1H, m, H-24) suggested the existence of secondary hydroxyl group. The ^1H -NMR spectrum showed the presence of six tertiary methyl groups at δ_{H} 0.75 (3H, s, Me-29), 0.88 (3H, s, Me-30), a sharp signal appearing at δ_{H} 0.97 (6H, s, Me-18, Me-28), 1.12 (3H, s, Me-26), 1.24 (3H, s, Me-27) and one methyl group at δ_{H} 0.87 (3H, d, $J = 3.2$ Hz, Me-

21) coupled with H-20 (methine proton) gave a doublet.

The ^{13}C -NMR spectrum (125 MHz, CDCl_3) [Table 1] of compound **3** showed the presence of 30 carbon signals. Determination of the multiplicity was carried out by DEPT experiments which revealed the presence of 7 methyl groups, 11 methylene groups, 6 methine groups with two oxygenated carbons at δ_{C} 78.83 for (C-3), 79.63 for (C-24) and 6 quaternary carbon signals with 1 oxygenated at δ_{C} 76.74 for (C-25).

The structure of compound **3** was deduced from the comparison of its spectral data with those of literature and was identified as cycloartan-3, 24, 25-triol.^[23-26]

ACKNOWLEDGMENT

The authors thank Dr. Olga Tzakou for providing the plant species.

REFERENCES

- Webster G. Systematic of the *Euphorbiaceae*: Introduction. *Ann Miss Bot Gard* 1994;81:133-44.
- Singla A, Pathak K. Phytoconstituents of *Euphorbia* species. *Fitoterapia* 1990;61:483-516.
- Appendino G, Spagliardi P, Ballero M, Seu G. Macrocyclic diterpenes from *Euphorbia hyberna* L. subsp. *insularis* and their reaction with oxyphilic reagents. *Fitoterapia* 2002;73:576-82.
- Haba H, Lavaud C, Harkat H, Alabdul Magid A, Marcourt L, Benkhaled M. Diterpenoids and triterpenoids from *Euphorbia guyoniana*. *Phytochemistry* 2007;68:1255-60.
- Hohmann J, Evanics F, Dombi G, Szabó P. Salicifoline and Salicinolide, new diterpene polyesters from *Euphorbia salicifolia*. *Tetrahedron Lett* 2001;42:6581-4.
- Evans FJ, Taylor SE. Proinflammatory tumor-promoting and anti-tumor diterpenes of the plant families Euphorbiaceae and Thymillaeaceae. In: Herz W, Grisebach H, Kirby GW, editors. *Progress in the Chemistry of Organic Natural Products*. Springer-Verlag, New York, 1983, pp. 1-99.
- Sahai R, Rastogi RP, Jakupovic J, Bohlmann F. A diterpene from *Euphorbia maddenii*. *Phytochemistry* 1981;20:1665-7.
- Uemura D, Katayama C, Uno E, Kansuine B. A novel multi oxygenated diterpene from *Euphorbia kansui* Liou. *Tetrahedron Lett* 1975;21:1703-6.
- Corea G, Fattorusso E, Lanzotti V, Tagliatalata-Scafati O, Appendino G, Ballero M, et al. Jatrophone diterpenes as P-glycoprotein inhibitors. First insights of structure-activity relationships and discovery of a new, powerful lead. *J Med Chem* 2003;46:3395-402.
- Abdelgaleil SA, Kassem SM, Doe M, Baba M, Nakatani M. Diterpenoids from *Euphorbia paralias*. *Phytochemistry* 2001;58:1135-9.
- Hohmann J, Evanics F, Dombi G, Molnár J, Szabó P. Euphosalicin, a new diterpene polyester with multidrug resistance reversing activity from *Euphorbia salicifolia*. *Tetrahedron* 2001;57:211-5.
- Zhou TX, Bao GH, Ma QG, Qin GW, Che CT, Liv Y, Wang C, Zheng QT, Langduin C. A novel dimeric diterpenoid from the roots of *Euphorbia fischeriana*. *Tetrahedron Lett* 2003;44:135-7.
- Hohmann J, Rédei D, Evanics F, Kálmán A, Argay G, Bartok T. Serrulatin A and B. New Diterpene Polyesters from *Euphorbia serrulata*. *Tetrahedron* 2000;56:3619-23.
- Ilyas M, Parveen M, Amin KM. Neriifolione. A triterpene from *Euphorbia nerifolia*. *Phytochemistry* 1998;48:561-3.
- Lima EM, Medeiros JM, Davin LB. Pentacyclic triterpenes from *Euphorbia stygiana*. *Phytochemistry* 2003;63:421-5.
- Shi JG, Shi YP, Jia, ZJ. Sesquiterpenoids from *Euphorbia wangii*. *Phytochemistry* 1997;45:343-7.
- Rizk AM, Youssef AM, Diab MA, Salem HM. Constituents of Egyptian Euphorbiaceae. Part 2: Flavonoids of *Euphorbia segetalis* Pharmazie 1976;31:405-7.
- Fatope MO, Zeng L, Ohayaga JE, Shi G, McLaughlin JL. Selectively cytotoxic diterpenes from *Euphorbia poissonii*. *J Med Chem* 1996;39:1005-8.
- Margret MO, Michel DB, Christopher SC, Barbara JWC. Betulin and lupeol in bark from four white-barked birches. *Phytochemistry* 1988;27:2175-6.
- Tinto WF, Blair LC, Alli A, Reynolds WF, Mclean S. Lupane triterpenoids of *Salacia cordata*. *J Nat Prod* 1992;55:395-8.
- Khan MT, Khan SB, Ather A. Tyrosinase inhibitory cycloartane type triterpenoids from the methanol extract of the whole plant of *Amberboa ramosa* Jafri and their structure-activity relationship. *Bioorg Med Chem* 2006;14:938-43.
- Garcez FR, Núñez CV, Garcez WS, Almeida RM, Roque NF. Sesquiterpenes, limonoid and coumarin from the wood bark of *Guarea guidonia*. *Planta Med* 1998;64:7-80.
- Akira I, shintaro O, Takako S, Hiroko M, Yuka I, Dedy D, Tsutomu N. Cycloartane triterpenoids from *Aglaia harmsiana*. *Phytochemistry* 1997;46:379-81.
- Hisham A, Bai A, JayaKumar G, Nair MS, Fujimoto Y. Triterpenoids from *Dysoxylum malabaricum*. *Phytochemistry* 2001;56:331-4.
- Nyemba AM, Mpondo TN, Connolly JD, Rycroft DS. Cycloartane derivatives from *Garcinia lucida*. *Phytochemistry* 1990;29:994-7.
- Satti NK, Suri OP, Dhar KL, Atal CK, Kawasaki T, Miyahara K, Kawano S. High resolution NMR and x-ray crystallography data of caudicifolin from *Euphorbia acaulis*. *Phytochemistry* 1986;25:1411-3.

Source of Support: Nil, Conflict of Interest: None declared.