

A Standardized Composition Comprised of Extracts from *Rosmarinus officinalis*, *Annona squamosa* and *Zanthoxylum clava-herculis* for Cellulite

Mesfin Yimam, Young-Chul Lee¹, Ping Jiao, Mei Hong, Lidia Brownell, Qi Jia

Unigen Inc., Seattle, WA 98121, USA, ¹Unigen Inc., Cheonan-Si, Chungnam 330-863, Korea

ABSTRACT

Background: Cellulite, characterized by changes in the skin morphology presented as dimpled or puckered skin appearance, is highly prevalent among postadolescent women. Cellulite management ranges from topical cream applications to invasive procedures. While some interventions showed improvements in physical appearances of affected areas, so far, none have reversed the condition to a full recovery. These unsuccessful measures signify the intricate nature of cellulite etiology highlighting its complexity leading to the possibility for a combination treatment approach to target multiple mechanisms. **Materials and Methods:** We screened our plant library for extracts that reduce cellular lipid accumulation, improve microcirculation, possess high total antioxidant capacity, significant anti-platelet aggregation, and anti-inflammatory activities using lipid accumulation assay in 3T3-L1 cells, Croton oil-induced hemorrhoid test in rats as a model for microcirculation, anti-platelet aggregation assay, nitric oxide (NO) inhibition assay, and 1,1-diphenyl-2-picrylhydrazyl assay.

Results: Three known botanicals such as *Rosemary officinalis*, *Annona squamosa* and *Zanthoxylum clava-herculis* were identified as lead extracts in these tests. Treatment of 3T3 cell with *A. squamosa* at 1 µg/ml resulted in 68.8% reduction in lipid accumulation. In croton oil-induced hemorrhoid study, *Z. clava-herculis* reduced the recto-anus coefficient by 79.6% at 6 mg/kg indicating improvement in microcirculations. Similarly, *R. officinalis* caused inhibition of 82%, 71.8%, and 91.8% in platelet aggregation, NO production and free radical generation at 31.25 µg/ml, 6.2 µg/ml, and 40 µg/ml concentrations suggesting its anti-oxidant, and anti-inflammatory activities. **Conclusions:** Data depicted here suggest that formulation of these well-known botanicals at a specific ratio perhaps may yield a composition with a much wider spectrum of mechanisms of actions to impact the multiple pathways involved in cellulite onset, continuation, or exacerbations.


Key words: *Annona squamosa*, cellulite, *Rosemary officinalis*, *Zanthoxylum clava-herculis*

SUMMARY

- Cellulite represents one of the main esthetic concerns of women with a likely cause of psychological insecurities. Its pathophysiology involves multiple pathways that include vascular, adipose tissues, inflammation, structural and physiological.
- Treatment strategies for cellulite comprises increasing microcirculation flow,

reducing lipogenesis, promoting lipolysis, free radicals scavenging or formation reduction, anti-inflammation and other invasive procedures.

- We screened our plant library for extracts that reduces cellular lipid accumulation, improves microcirculation, possesses high total antioxidant capacity, inhibits platelet aggregation, and moderates inflammation.
- Botanical extracts from *Rosmarinus officinalis*, *Annona squamosa* and *Zanthoxylum clava-herculis* were identified as leads and formulated to yield a standardized composition designated as UP1307 and suggested its usage for cellulite.



Abbreviations Used: GMP: Good Manufacturing Practice; CA: Carnosic acid; NF-kB: nuclear factor-kB; HPLC: high-performance liquid chromatography; EtOH: Ethanol; DMEM: Dulbecco's modified Eagle's medium; FBS: fetal bovine serum; SD: Sprague Dawley; RAC: recto-anus coefficient; LPS: Lipopolysaccharide; DPPH: 1,1-diphenyl-2-picryl-hydrazyl; TNF-α: tumor necrosis factor; NO: Nitric oxide

Correspondence:

Dr. Mesfin Yimam,
Unigen Inc., 3005 1st Ave, Seattle, WA 98121,
USA.

E-mail: myimam@unigen.net

DOI: 10.4103/pr.pr_70_17

Access this article online

Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

Cellulite, characterized by changes in the skin morphology presented as dimpled or puckered skin appearance around the posterolateral region, is highly prevalent among women. Records suggest that about 85% of post-adolescent women have some degree of cellulite.^[1] It represents one of the main esthetic concerns of women with a likely cause of psychological insecurities that may lead to a desperate need for cosmetic or treatment interventions.

Well-defined specific etiology of cellulite has remained elusive for decades. Through time, substantial speculations have been reported describing the complex nature and the tangled co-presence of multiple factors involving in the initiation, perpetuation, and exacerbation of

the condition rendering multiple definitions.^[2-5] It is a multifactorial condition where its occurrence believed to be associated with mainly

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Yimam M, Lee YC, Jiao P, Hong M, Brownell L, Jia Q. A standardized composition comprised of extracts from *Rosmarinus officinalis*, *Annona squamosa* and *Zanthoxylum clava-herculis* for cellulite. Phcog Res 2017;9:319-24.

(1) vascular-deficiencies in lymphatic drainage and microvascular circulation leading to edema within the subcutaneous tissue,^[6,7] (2) abnormal adipose deposition-localized abnormal fat deposition in the subcutaneous tissue (3) inflammation-chronic inflammatory process (4) structural – gender dependent architectural differences in subcutaneous tissues and changes in connective tissues, and (5) physiologic – a normal physiologic state that maximizes adipose retention to ensure adequate caloric availability for pregnancy and lactation in postadolescent women.^[2]

In spite of the limited understanding in the pathophysiology of cellulite, significant treatment strategies such as (a) increasing the microcirculation flow, (b) reducing lipogenesis and promoting lipolysis, (c) preventing the free radical formation or scavenging free radicals, and (d) modulating with anti-inflammatory and anti-edematous agents have been proposed. Varieties of formulations have been currently dispensed for cellulite. Some of these preparations contain methylxanthines (caffeine, aminophylline, theophylline, and theobromine), retinol (Vitamin A), alpha-tocopherol (Vitamin E), and ascorbic acid (Vitamin C) as the major active constituents. Extracts from *Ginkgo biloba*, *Centella asiatica*, *Ruscus aculeatus*, *Ananas sativus*, *Chenopodium quinoa* seed, and Yuzu seed are also available for a similar indication. However while some interventions showed improvements in physical appearances of affected areas, so far, none have reversed the condition to a full recovery. These unsuccessful measures further signify the intricate nature of the condition highlighting its complexity leading to the discussion of a combination therapy to target multiple mechanisms. In this regard, we screened our plant library for extracts that reduce cellular lipid accumulation, possess the high total antioxidant capacity, significant anti-platelet aggregation and anti-inflammatory activities. Three known botanicals such as *Rosemary officinalis* (RM504-C), *Annona squamosa* (RM606) and *Zanthoxylum clava-herculis* (UP342-C) were identified in these tests as leads and formulated at a specific ratio to yield a composition designated as UP1307.

Significant reports have been documented describing the various traditional and contemporary ethnopharmacological applications of these three medicinal plants. For example, *A. squamosa* L. (family: *Annonaceae*), also known as sugar apple, or custard apple, is the most widely grown *Annona* fruit tree in tropical and subtropical areas comprising of about 130 genera and more than 2300 species. The plant is a native of the West Indies and is currently cultivated throughout India mainly for its edible fruit.^[8,9] This plant is widely known to possess several medicinal properties^[10,11] with various phytochemical and biological activities.^[12,13] Partially purified flavonoids of aqueous *Annona* leaves extract to possess antimicrobial and insecticidal activity.^[14] Annonaceous acetogenins constitute a series of natural products isolated exclusively from *Annonaceae* plants with a broad range of biological activities such as anticancer, cytotoxic, antiparasitic, pesticidal, and immunosuppressive activities.^[8,9]

Rosmarinus officinalis L. is an evergreen perennial shrub native to Europe and cultivated in many parts of the world. Rosemary leaves are often used as spices and flavoring agents because of the desirable flavor and antioxidant activity of the dried leaves of the plant. There is an increasing interest in the pharmaceutical properties of rosemary, being used in conventional medicine to improve memory and relieve pain, for its antimicrobial, hepatoprotective, anti-inflammatory, anti-tumorigenic, or chemo-preventive activities.^[15] The major bioactive component of rosemary extract, carnosol and carnosic acid (CA) have showed a wide range of activities, which include adipogenesis inhibition^[16,17] glucose and lipid metabolism regulation,^[18] weight gain reduction,^[19,20] cholesterol levels and glycemia improvement,^[20] liver steatosis improvement^[19] and gastric lipase inhibition yielding body weight and plasma lipids

management.^[21] CA also inhibits cytokine-induced adhesion molecules expression and monocyte adhesion to endothelial cells through a mechanism that involves nuclear factor- κ B (NF- κ B), which could be involved in the anti-inflammatory properties of this compound. The CA may undergo an oxidative degradation and rearrangement cascade, giving rise to other rosemary antioxidant compounds such as carnosol, rosmanol, galdosol, and rosmariquinone.^[22]

Similarly, *Zanthoxylum clava-herculis* L. and *Zanthoxylum americanum* Mill, commonly known as Southern and Northern prickly ash, respectively, belong to the Yellow Wood (Rutaceae) family and has a long historical use as a botanical remedy. *Zanthoxylum* extract also referred as the toothache tree, conventionally been used by the Native American as a remedy for toothaches where the root-bark and berries having been listed in the old version of the United States Pharmacopoeia. The bark is greatly recommended for chronic rheumatism, typhoid, ulcers, colic, skin sores, diarrhea, indigestion, and circulatory problems.^[23,24] The bark of Prickly Ash contains aporphine alkaloids (magnoflorine, laurifoline, and liriodenine), benzo-phenanthridine alkaloids (chelerythrine), lignan (asarinin), coumarins (xanthoxyletin and xanthyletin), long chain amides (herculin), and essential oil.^[25]

In the present study, we evaluated each component of the composition on suggested key mechanisms of action of cellulite (such as adipogenesis, vasorelaxation, platelet aggregation, inflammation, and free radical formation) and recommended further investigation on their composition for the potential use in cellulite.

MATERIALS AND METHODS

Materials

Annona squamosa L

The sugar apple fruit extract is manufactured through a series of proprietary processing steps which have been validated and performed in accordance with Good Manufacturing Practice (GMP). *Annona* dried whole fruit was ground to a particle size of no larger than 2 ml. Dried ground sugar apple fruit (100 kg) was then transferred to a supercritical fluid extractor (pressure 25 Mpa, temperature 50°C, CO₂ flow 70 ml/min, 2 h) and ethyl alcohol (95% food grade ethanol, quantity is 30% of the raw material) was added. The supercritical fluid CO₂ extract was evaporated under vacuum. The extraction yield is 3.7%. The contents of two biomarkers were quantified by high-performance liquid chromatography (HPLC) method with Squamocin not <1%, and Kaurenoic acid content not <4% in the extract. Squamocin and Kaurenoic Acid in the *A. squamosa* fruit extract were quantified with a Luna C-18 reversed-phase column (Phenomenex, 10 μ m, 250 mm \times 4.6 mm) in an Agilent HPLC system at 210 nm.

Zanthoxylum clava-herculis

A total of 800 kg of fresh *Z. clava-herculis* L. (Prickly Ash) tree bark was dried, cut, crushed, and then extracted with approximately 3-fold volume (2400 L) of 90% ethyl alcohol in water (v/v); the extraction was carried on at 90°C for 8 h. The ethanol solution was filtered to obtain the supernatant which was then concentrated with an evaporator under vacuum at 40°C. This extraction and concentration procedure was repeated three times. The extraction solutions were then combined and concentrated together. The concentrated solution stayed for 24 h in the refrigerator to obtain a supernatant. The supernatant was vacuum-dried to obtain 87.5 kg of Prickly Ash EtOH extract powder. The extraction yield was about 10.9% (w/w). Two biomarkers-Magnoflorine and Laurifoline in Prickly Ash extract were quantified with a Luna C18 reversed-phase column (Phenomenex, 5 μ m, 250 mm \times 4.6 mm) in an Agilent 1200 HPLC system at 275 nm.

Rosemary officinalis

Dried 29 kg of Rosemary leaf was extracted with approximately 10-fold volume (300 L) of 95% ethyl alcohol at 40°C. Above-described procedure was repeated two times. The extraction solutions were concentrated until the volume became 1/30 the original volumes. The concentrated solutions were then combined and evaporated again to reduce the volume of concentrated solution until 1/90 the volume of the original extraction solution was achieved. The concentrated solution was left at room temperature for 24 h to allow separation into a supernatant and precipitate-layer. Concentrated supernatant was dried by vacuum drying to obtain rosemary extract that contained between 30% and 60% CA. CA was quantified with a Luna C18 reversed-phase column (Phenomenex, 5 µm, 250 mm × 4.6 mm) in an Agilent HPLC system at 206 nm.

Lipid accumulation assay in 3T3-L1 cells

3T3 L1 mouse embryo fibroblasts were purchased from American Type Culture Collection. Cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (GIBCO) containing 10% bovine calf serum until confluent. Two days after postconfluence (D0), the cells were stimulated to differentiate with DMEM containing 10% fetal bovine serum (FBS), 5 µg/ml insulin, 0.5 mM 3-isobutyl-1-methylxanthine, and 1 µM dexamethasone for 2 days (D2). Cells were then maintained in 10% FBS/DMEM medium with 5 µg/ml insulin for another 2 days (D4), followed by culturing with 10% FBS/DMEM medium for 4 days (D8). Test samples were treated from day 0–8 of adipogenesis with RM606 at 0.0625 µg/ml, 0.125 µg/ml, 0.25 µg/ml, 0.35 µg/ml, 0.5 µg/ml, and 1 µg/ml concentrations, and medium was changed every 2 days. On day 8, lipid droplets on cells were stained with Oil Red-O and measured at 510 nm. Tumor necrosis factor-α (TNF-α) at a concentration of 10 ng/ml was used as a positive control.

Croton oil induced haemorrhoid

Male SD rats (6 weeks old, approximately 140 g) were purchased from Harlan Sprague–Dawley and allowed to acclimate for 1 week. The rats were maintained in a pathogen-free facility in accordance with the National Research Council of Laboratory Animal care and use guidelines. Each experiment was performed with age-matched rats 7–8 weeks old. The croton oil-induced hemorrhoid model in rats was performed according to the method published by Nishiki *et al.*^[26] In brief, a cotton swab with a diameter of 4 mm soaked with 0.16 mL of inducer (deionized water: Pyridine: Ethyl ether: 6% croton oil/ethyl ether (1:4:5:10) was applied to the rat's anus (rectoanal portion, 20 mm from anal opening) for 12 s. The final concentration of croton oil was 3%. Rats were administered with UP342-C suspended in saline at oral doses of 0.67 mg/kg, 2.0 mg/kg and 6.0 mg/kg 24 h after hemorrhoid induction. Twenty-four hours later, recto-anus tissue (approx. 10 mm long) was isolated after the rats were euthanized. The weights of rat body and recto-anus were measured. The recto-anus coefficient (RAC) was calculated using the formula: weight of recto-anus (mg)/body weight (g).

Anti-platelet aggregation assay

Whole blood was collected from male Sprague–Dawley rats and then transferred into 15 mL test tube containing 1 mL of anticoagulant citrate/dextrose solution (ACD, 85 mM trisodium citrate, 83 mM dextrose, and 21 mM citric acid). Blood was centrifuged at 170 ×g for 7 min. to obtain platelet-rich plasma, which was further centrifuged at 120 ×g for 7 min to remove residual erythrocytes. This platelet-rich plasma was centrifuged twice more at 350 ×g with a washing buffer for 10 min to remove the ACD solution, and then platelet precipitates were adjusted to (3 × 10⁸/mL) for aggregation assay in Tyrode buffer (137

mM of NaCl, 12 mM of NaHCO₃, 5.5 mM of glucose, 2 mM of KCl, 1 mM of MgCl₂, 0.3 mM of NaHPO₄, and pH 7.4). The washed platelets were preincubated at 37°C for 2 min with RM504-C or vehicle (<0.1%) and then stimulated with collagen. Collagen is a well-known soluble agonist for platelet aggregation and thrombus formation. The reaction mixture was further incubated for 5 min with stirring at 170 ×g. Aggregation was monitored by measuring light transmission in an aggregometer (Chronolog, Havertown, PA, USA).

Nitric oxide inhibition assay

Overproduction of NO in macrophages is a hallmark of inflammation. Therefore, we determined the effect of RM504-C (6.25–100 µg/ml) in LPS (0.1 µg/ml) stimulated RAW264.7 cells. RAW264.7 (Korean Cell Line Bank, South Korea) were cultured and maintained in DMEM (Dalseogu, South Korea) enriched with 10% heat-inactivated FBS (WelGene Co., South Korea), 100 µg/ml streptomycin, and 100 U/ml penicillin (Lonza, MD, USA) in a humidified atmosphere of 5% CO₂ at 37°C. Cultured RAW264.7 cells (4 × 10⁵) were pretreated with or without SCME (1,3,10 µg/ml) for 30 min and then stimulated with lipopolysaccharide (LPS, 0.1 µg/ml) for 18 h in a 96 well plate. The cell culture supernatant (100 µl) was mixed with an equal volume of Griess reagent (1% sulphanilamide in 5% phosphoric acid (H₃PO₄) and 0.1% N-1-naphthylenediamine dihydrochloride in deionized distilled water) and then subjected for plate reading at 540 nm in an ELISA reader. The accumulated nitrite in the culture medium was quantified using a standard of NaNO₂ (1 mM).

1,1-Diphenyl-2-picrylhydrazyl assay

The stable free radical 1,1-diphenyl-2-picryl-hydrazyl (DPPH) was used to examine the free radical scavenging activity of Rosemary (RM504-C) extract samples. In brief, a 0.2 mM solution of DPPH was prepared in DMSO, which was mixed with each extract sample at concentrations of 1, 5, 10, 20, and 40 µg/ml. After a 30 min incubation in the dark, absorbance was measured at 517 nm with a spectrophotometer. A decrease in solution absorbance indicates a decrease of DPPH and an anti-oxidant effect. The anti-oxidant activity is expressed as a percent inhibition.

RESULTS

Effect of *Annona squamosa* extract (RM606) on lipid accumulation

Dose correlated reduction in cellular lipid accumulation was observed when 3T3 cells were treated with RM606 at 0.0625 µg/ml to 1 µg/ml concentration. Compared to control, percent inhibition of 32.1%, 45.6%, and 68.8% were observed for RM606 at 0.35, 0.5, and 1 µg/ml, respectively [Figure 1] with an IC₅₀ value as 0.489 µg/ml. The positive control, TNF-α showed 77.1% inhibition compared to control.

Effect of *Zanthoxylum clava-herculis* (UP342-C) on microcirculation improvement

The edema developed linearly until 7–8 h after application of croton oil and the severity of the edema was sustained for more than 24 h. Vehicle alone did cause significant swelling of the recto-anus as compared with the rats without the extract [Figure 2]. The average RAC of vehicle-treated croton oil-induced rats was 1.66 which was 40.7% higher than that of the normal control (RAC = 1.18) [Figure 2]. However, treatment with UP342-C extract reduced the swelling of the recto-anus by 79.5%, 67.5%, and 42.1%, for dosages 6 mg/kg, 2 mg/kg, and 0.67 mg/kg, respectively [Figure 2]. The same preparation was tested at 6 mg/kg for topical application and found a 93.1% RAC inhibition with statistical significance when compared to vehicle-treated croton oil-induced rats with *P* = 0.02 (USP NO. 6210680B1) indicating its effect in microcirculation.




Figure 1: Effect of *Annona squamosa* extract (RM606) on lipid accumulation. 3T3 L1 cells were treated with RM606 at 0.0625 µg/ml, 0.125 µg/ml, 0.25 µg/ml, 0.35 µg/ml, 0.5 µg/ml, and 1 µg/ml concentrations. Lipid droplets on cells were stained with Oil Red-O and measured at 510 nm on day 8. Data are expressed as percent inhibition of lipid accumulation




Figure 3: Anti-platelet aggregation effects of *Rosmarinus officinalis* (RM504-C). Platelets were pre-incubated at 37°C for 2 min with RM504-C or vehicle (<0.1%) and then stimulated with collagen. The reaction mixture was incubated for 5 min. with stirring at 170 ×g. Aggregation was monitored by measuring light transmission in an aggregometer. Data are expressed as percent inhibition of aggregation

Anti-platelet aggregation effects of *Rosemary officinalis* (RM504-C)

CA enriched rosemary extract (RM404-C) showed a dose-dependent inhibition of collagen-induced platelet aggregation in rat platelets [Figure 3]. The highest inhibition, 82%, in platelet aggregation was observed when RM404-C was used at 31.25 µg/ml concentration.

Nitric oxide inhibition effects of rosemary extract (RM504-C)

Nitric oxide (NO) overproduction can serve as proinflammatory mediator at pathologic conditions to induce and/or exacerbate inflammation. Data in Figure 4 showed that RM504-C was a potent inhibitor of NO production suggesting its anti-inflammatory activity. Dose-dependent inhibition of




Figure 2: Effect of *Zanthoxylum clava-herculis* (UP342-C) on microcirculation improvement. Croton oil-induced hemorrhoid in rats was used as a disease model. Rats were administered with UP342-C suspended in Saline at oral doses of 0.67 mg/kg, 2.0 mg/kg, and 6.0 mg/kg 24 h after hemorrhoid induction. Control group is for the vehicle control (Saline). The recto-anus coefficient was determined 24 h posttreatment. Recto-anus coefficient was calculated using the formula: Weight of recto-anus (mg)/body weight (g)




Figure 4: Nitric oxide inhibition effects of rosemary extract (RM504-C). Lipopolysaccharide (0.1 µg/ml) stimulated RAW264.7 cells were used to determine the over production of nitric oxide. Controls are unstimulated cells expected to have minimal to no production of nitric oxide. Cells were treated with RM504-C at 6.25–100 µg/ml concentrations. Data expressed as mean ± standard deviation of triplicates

NO was observed in this assay. At concentrations as low as 6.2 µg/ml, a 71.8% inhibition in production of NO was observed. Complete inhibition in NO production was noted for concentrations above 12.5 µg/ml. In the unstimulated control group, no production of NO was observed. Controls are unstimulated cells expected to have very low level or no radical production.

Anti-oxidant effect of rosemary extract (RM504-C)

DPPH assay is one of the most frequently used assays to determine the anti-oxidant activity of compounds. CA enriched rosemary extract (RM404-C) showed a dose-dependent inhibition in generation of free radicals as measured by DDPH assay. Inhibitions of 0.5%, 13.6%,




Figure 5: Anti-oxidant effect of rosemary extract (RM504-C) in 1,1-diphenyl-2-picrylhydrazyl assay. The extract was tested at concentrations of 1, 5, 10, 20 and 40 µg/ml. The anti-oxidant activity is expressed as a percent inhibition

38.4%, 85.2%, and 91.8% were observed for RM504-C at 1, 5, 10, 20, and 40 µg/ml, respectively [Figure 5].

DISCUSSION

The present study depicts preclinical data from cellular lipid accumulation, microcirculation improvement, anti-platelet aggregation, NO inhibition, and DPPH assays for three well-known medicinal plant extracts such as *R. officinalis*, *A. squamosa* and *Z. clava-herculis*. The significance of these findings in association with cellulite have been discussed.

Cellulite is a nonpathologic skin condition mainly known for its negative impacts on quality of life with no mortality or morbidity. However, it takes a serious psychological toll on affected women. Blood and lymphatic microcirculation damage are believed to give rise to structural changes in the fatty layer and surrounding collagen matrix at cellulite pathophysiology. Subsequently, subcutaneous edema could also appear as a sequel as a result of superficial microcirculation insufficiency due to altered permeability of blood vessels. Cellulite management ranges from topical cream applications to invasive procedures, including laser-assisted lipolysis and liposuction. Despite the significant progress in the field of combating cellulite, currently, there are no treatment options successfully curing the underline causes. Among the various predisposing factors that influence cellulite, both microvascular and lymphatic circulation impairment are considered the primary etiological factors. As a result, significant numbers of products are commercially available in various formulations targeting microcirculation. For example, *C. asiatica* extracts for microcirculation and an anti-inflammation,^[27] *R. aculeatus* for improving lymphatic drainage,^[28] *Carica papaya* as anti-edema,^[29] and Common ivy (*Hedera helix*) for lymphatic drainage^[30] are some of the topically applicable botanical extracts with limited success in treating cellulite. These materials believed to impose their desired outcome by reducing blood viscosity, inhibiting platelet aggregation, moderating vascular permeability, and improving vascular tone, which may lead to improved microcirculation. Aligning with these proposed underlying mechanisms, in the present study, significant inhibition in platelet aggregation (as a result of *R. officinalis*) and improved microcirculation by *Zanthoxylum clavaherculis* were observed for constituents of the composition suggesting their beneficial usage for improved microcirculation. Recently, it has also been shown that the major bioactive component of *Zanthoxylum* extract, Magnoflorine,

to have a dual target effect in NF-κB inhibition and β₂-adrenoceptor agonist activities both of which have a primary contribution in the pathophysiology of cellulite.^[31] While the former activity modulates the inflammatory pathway, the later may lead to smooth muscle relaxation and hence vasodilation which could have a measurable impact in improving microcirculation rendering their application in cellulite management.

Asserting the multifactorial causative agents of cellulite, adipogenesis, and lipolysis play a major role at the cellulite-prone areas in the pathophysiology of cellulite. Physiologic and biochemical properties of adipose tissue in these areas have been shown to be different from subcutaneous adipose tissue elsewhere in both hormonal responsiveness and metabolic activities. Among the few neurotransmitters that affect the acute lipolysis of adipose tissues (e.g., catecholamine) work through the activation of the beta-adrenergic receptor.^[32,33] Stimulation of these receptors as an agonist has a positive effect on lipolysis. For instance, β-agonists such as methylxanthines (e.g., caffeine and aminophylline) preparation are widely available on the market for topical use for cellulite believed to impose their activity through acting directly on adipose cells, promoting lipolysis, inhibiting phosphodiesterase, increasing levels of cyclic adenosine monophosphate and/or activating the triglyceride lipase enzyme and breaks down triglycerides into free acids and glycerol.^[34,35] Here again, the β₂-adrenoceptor agonist alkaloid extract from *Zanthoxylum* (e.g., magnoflorine) may share similar mechanisms in promoting lipolysis situating the extract with additional benefit in cellulite intervention. Most importantly, in our study, extract from *A. squamosa* showed a direct impact on adipogenesis as demonstrated by the significant reduction in lipid accumulation in 3T3 cells. In fact, during our discovery process, we screened more than 5300 plants extracts using the fat cells isolated from the human adipose tissues and identified acetogenins as the primary actives from *A. squamosa* with an IC₅₀ of 1 ng/ml in inhibition of adipogenesis. Substantiating our findings, Gokaraju *et al.* have reported the effect of the major active component of Squamosa extract, acetogenins, in down regulating adipogenesis differentiating markers and perilipin protein revealing its inhibiting adipogenesis and/or accelerating lipolysis activities reinforcing its usage in cellulite modulation.^[36]

There are reasonable assumptions for the presence of inflammatory processes and hence free radical generation or vice versa at the time of deposition of dystrophic adipose tissues during the pathogenesis of cellulite.^[37] For example, abnormal changes in the endothelium and edema formation are the two possible pathological signals suggestive of the existence of these phenomena. Moreover, the isolation of macrophages and monocytes from the biopsies of fibrous septae of patients with cellulite were further considered as an indication of involvement of chronic low-grade inflammation in the pathophysiology of cellulite.^[38] In these regards, extracts from all the three components of the composition, *Rosmarinus officinalis*,^[39] *A. squamosa*,^[40] and *Xanthoxylum americanum*^[41] have been previously reported to possess anti-inflammatory activities that strengthen their usage in cellulite intervention. In particular, the free radical scavenging and inhibition of NO activities of *Rosmarinus officinalis* have an extract that been demonstrated in our assays attest its strong anti-oxidant and anti-inflammatory activities. These findings further fortify the argument that each component of the composition has a significant share in addressing one or multiple pathways among the suggested possible mechanisms of action in cellulite pathology. Some topical formulations composed of *G. biloba* as anti-inflammatory and anti-oxidant,^[1] Red grapes (*Vitis vinifera*) as an antioxidant,^[30] and *Carica papaya* as anti-inflammatory^[29] are examples of plant extracts with a similar mechanism of actions indicated for cellulite.

The current study is not without a limitation. While statistically significant results were reported in all the disclosed assays for each extract tested, for some, no positive controls were used. As a result, the relative potency of the extracts cannot be determined against known standard references.

CONCLUSIONS

Collectively, botanical formulations applied topically as anti-cellulite intend to reduce lipogenesis, activate lipolysis, restore subcutaneous tissue normal structure, reduce free radical generation or scavenge free radicals, prevent inflammation, increase microcirculation, and lymphatic drainage. Data depicted here suggest that formulation of these well-known botanicals at specific ratio perhaps may yield a composition with a much wider spectrum of mechanisms of actions to impact the multiple pathways involved in cellulite onset, continuation or exacerbations. Hence, we suggest further mechanism based elucidation of their composition for cellulite management and validation of hypothesis in human clinical study.

Acknowledgments

The authors would like to express their best gratitude to Dr. Eujin Hyun, Dr. Edward, Dr. Wenwen Ma, and Dr. Min Chu, and Unigen team for their incalculable support for the completion of this research.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Hexsel D, Orlandi C, Zechmeister do Prado D. Botanical extracts used in the treatment of cellulite. *Dermatol Surg* 2005;31:866-72.
- Piérard GE. Commentary on cellulite: Skin mechanobiology and the waist-to-hip ratio. *J Cosmet Dermatol* 2005;4:151-2.
- Curri SB, Ryan TJ. Panniculopathy and fibrosclerosis of the female breast and thigh. *Clin Dermatol* 1989;7:107-19.
- Rossi AB, Vergnanini AL. Cellulite: A review. *J Eur Acad Dermatol Venerol* 2000;14:251-62.
- Goldman MP. Cellulite: A review of current treatments. *Cosmet Dermatol* 2002;15:17-20.
- Cellulite meltdown? *Harv Womens Health Watch* 1998;5:7.
- Piérard GE, Nizet JL, Piérard-Franchimont C. Cellulite: From standing fat herniation to hypodermal stretch marks. *Am J Dermatopathol* 2000;22:34-7.
- Rupprecht JK, Hui YH, McLaughlin JL. Annonaceous acetogenins: A review. *J Nat Prod* 1990;53:237-78.
- Shirwaikar A, Rajendran K, Dinesh Kumar C, Bodla R. Antidiabetic activity of aqueous leaf extract of *Annona squamosa* in streptozotocin-nicotinamide type 2 diabetic rats. *J Ethnopharmacol* 2004;91:171-5.
- Yang H, Li X, Tang Y, Zhang N, Chen J, Cai B, *et al.* Supercritical fluid CO₂ extraction and simultaneous determination of eight annonaceous acetogenins in *Annona* genus plant seeds by HPLC-DAD method. *J Pharm Biomed Anal* 2009;49:140-4.
- Asolkar LV, Kakkar KK, Chakre OJ. Glossary of Indian Medicinal Plants with Active Principles. New Delhi: Publication and Information Directorate; 1992. p. 72-3.
- Vohora SB, Kumar I, Naqvi SA. Phytochemical, pharmacological, antibacterial and anti-ovulatory studies on *Annona squamosa*. *Planta Med* 1975;28:97-100.
- Lieb F, Nonfon M, Wachendorff-Neumann U, Wendisch D. Annonacins and annonastatin from *Annona squamosa*. *Planta Med* 1990;56:317-9.
- Kotkar HM, Mendki PS, Sadan SV, Jha SR, Upasani SM, Maheshwari VL, *et al.* Antimicrobial and pesticidal activity of partially purified flavonoids of *Annona squamosa*. *Pest Manag Sci* 2002;58:33-7.
- Hanson JR. Rosemary, the beneficial chemistry of a garden herb. *Sci Prog* 2016;99:83-91.
- Gaya M, Repetto V, Toneatto J, Anesini C, Piwien-Pilipuk G, Moreno S, *et al.* Antiadipogenic effect of carnolic acid, a natural compound present in *Rosmarinus officinalis*, is exerted through the C/EBPs and PPAR γ pathways at the onset of the differentiation program. *Biochim Biophys Acta* 2013;1830:3796-806.
- Takahashi T, Tabuchi T, Tamaki Y, Kosaka K, Takikawa Y, Satoh T, *et al.* Carnolic acid and carnolic acid inhibit adipocyte differentiation in mouse 3T3-L1 cells through induction of phase2 enzymes and activation of glutathione metabolism. *Biochem Biophys Res Commun* 2009;382:549-54.
- Tu Z, Moss-Pierce T, Ford P, Jiang TA. Rosemary (*Rosmarinus officinalis* L.) extract regulates glucose and lipid metabolism by activating AMPK and PPAR pathways in hepG2 cells. *J Agric Food Chem* 2013;61:2803-10.
- Harach T, Aprikian O, Monnard I, Moulin J, Membrez M, Béolor JC, *et al.* Rosemary (*Rosmarinus officinalis* L.) leaf extract limits weight gain and liver steatosis in mice fed a high-fat diet. *Planta Med* 2010;76:566-71.
- Ibarra A, Cases J, Roller M, Chiralt-Boix A, Coussaert A, Ripoll C, *et al.* Carnolic acid-rich rosemary (*Rosmarinus officinalis* L.) leaf extract limits weight gain and improves cholesterol levels and glycaemia in mice on a high-fat diet. *Br J Nutr* 2011;106:1182-9.
- Romo-Vaquero M, Selma MV, Larrosa M, Obiol M, García-Villalba R, González-Barrio R, *et al.* A rosemary extract rich in carnolic acid selectively modulates caecum microbiota and inhibits β -glucosidase activity, altering fiber and short chain fatty acids fecal excretion in lean and obese female rats. *PLoS One* 2014;9:e94687.
- López-Jiménez A, García-Caballero M, Medina MÁ, Quesada AR. Anti-angiogenic properties of carnolic acid and carnolic acid, two major dietary compounds from rosemary. *Eur J Nutr* 2013;52:85-95.
- Chevallier A. The Encyclopedia of Medicinal Plants. London, U. K: Dorling Kindersley Publications Inc.; 1996. p. 151.
- Bown D. Encyclopaedia of Herbs and their Uses. London: Dorling Kindersley; 1995. p. 372.
- Fish F, Waterman PG. Proceedings: Alkaloids in the bark of *Zanthoxylum clava-herculis*. *J Pharm Pharmacol* 1973;25 Suppl:115-6.
- Nishiki K, Nishinaga K, Kudoh D, Iwai K. Croton oil-induced hemorrhoid model in rat: Comparison of anti-inflammatory activity of diflucortolone valerate with other glucocorticoids. *Nihon Yakurigaku Zasshi* 1988;92:215-25.
- Hexsel D, Zechmeister do Prado D, Goldman MP. Topical management of cellulite. In: Goldman M, Hexsel D, editors. *Cellulite: Pathophysiology and Treatment*. London, England: Informa Health Care; 2010. p. 62-8.
- Rubanyi G, Marcelon G, Vanhoutte PM. Effect of temperature on the responsiveness of cutaneous veins to the extract of *Ruscus aculeatus*. *Gen Pharmacol* 1984;15:431-4.
- Carle R. Anti-inflammatory and spasmolytic botanical drugs. *Breast J Phytother* 1990;1:32-9.
- Facino RM, Carini M, Stefani R, Aldini G, Saibene L. Anti-elastase and anti-hyaluronidase activities of saponins and sapogenins from *Hedera helix*, *Aesculus hippocastanum*, and *Ruscus aculeatus*: Factors contributing to their efficacy in the treatment of venous insufficiency. *Arch Pharm (Weinheim)* 1995;328:720-4.
- Sun D, Han Y, Wang W, Wang Z, Ma X, Hou Y, *et al.* Screening and identification of *Caulis sinomenii* bioactive ingredients with dual-target NF- κ B inhibition and β -2-AR agonizing activities. *Biomed Chromatogr* 2016;30:1843-53.
- Wahrenberg H, Lönnqvist F, Arner P. Mechanisms underlying regional differences in lipolysis in human adipose tissue. *J Clin Invest* 1989;84:458-67.
- Leibel RL, Hirsch J. Site- and sex-related differences in adrenoreceptor status of human adipose tissue. *J Clin Endocrinol Metab* 1987;64:1205-10.
- Bertin C, Zunino H, Pittet JC, Beau P, Pineau P, Massonneau M, *et al.* A double-blind evaluation of the activity of an anti-cellulite product containing retinol, caffeine, and ruscogenine by a combination of several non-invasive methods. *J Cosmet Sci* 2001;52:199-210.
- Velasco MV, Tano CT, Machado-Santelli GM, Consiglieri VO, Kaneko TM, Baby AR, *et al.* Effects of caffeine and siloxanetriol alginate caffeine, as anticellulite agents, on fatty tissue: Histological evaluation. *J Cosmet Dermatol* 2008;7:23-9.
- Gokaraju GR, Gokaraju RR, Kanaka V, Gokaraju RR, Golakoti T, Sengupta K, *et al.* Extracts, Fractions and Compositions Comprising Acetogenins and their Applications. USPTO: US20120201884 A1; 2012.
- Terranova F, Berardesca E, Maibach H. Cellulite: Nature and aetiopathogenesis. *Int J Cosmet Sci* 2006;28:157-67.
- Kligman AM. Cellulite: Facts and fiction. *J Geriatr Dermatol* 1997;5:136-9.
- Ghasemzadeh Rahbardi M, Amin B, Mehri S, Mirnajafi-Zadeh SJ, Hosseinzadeh H. Anti-inflammatory effects of ethanolic extract of *Rosmarinus officinalis* L. and rosmarinic acid in a rat model of neuropathic pain. *Biomed Pharmacother* 2017;86:441-9.
- Wu P, Wu M, Xu L, Xie H, Wei X. Anti-inflammatory cyclopeptides from exocarps of sugar-apples. *Food Chem* 2014;152:23-8.
- Oriowo MA. Anti-inflammatory activity of piperonyl-4-acrylic isobutyl amide, an extractive from *Zanthoxylum zanthoxyloides*. *Planta Med* 1982;44:54-6.